

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/705, C12N 15/12, 15/63, 15/70, 15/79</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/48921</b> <b>(43) International Publication Date:</b> 30 September 1999 (30.09.99)
<b>(21) International Application Number:</b> PCT/US99/06573 <b>(22) International Filing Date:</b> 25 March 1999 (25.03.99)  <b>(30) Priority Data:</b> 60/079,501      26 March 1998 (26.03.98)      US  <b>(71) Applicants (for all designated States except US):</b> THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY [US/US]; Suite 350, 900 Welch Road, Palo Alto, CA 94304 (US). N.V. ORGANON [NL/NL]; Weth, Van Eschstraat 1, P.O. Box 20, NL-5340 BH Oss (NL).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> HSUEH, Aaron, J., W. [US/US]; 25 Ryan Court, Stanford, CA 94305 (US). HSU, Sheau, Yu [-/US]; Apartment 40, 234 Escuela Avenue, Mountain View, CA 94040 (US). LIANG, Shan-Guang [CN/US]; 438 Vennura Avenue #9, Palo Alto, CA 94306 (US). VAN DER SPEK, Petrus, Johannes [NL/NL]; Bremlaan 10, NL-5342 HM Oss (NL).  <b>(74) Agent:</b> FIELD, Bret, E.; Bozicevic, Field & Francis LLP, Suite 200, 285 Hamilton Avenue, Palo Alto, CA 94301 (US).		<b>(81) Designated States:</b> JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> NOVEL MAMMALIAN G-PROTEIN COUPLED RECEPTORS HAVING EXTRACELLULAR LEUCINE RICH REPEAT REGIONS  <b>(57) Abstract</b>  Isolated nucleotide compositions and sequences are provided for LGR4, LGR5 and LGR7 genes. The nucleic acid compositions find use in identifying homologous or related genes; in identifying endogenous ligands for these receptors; in producing compositions that modulate the expression or function of its encoded protein; for gene therapy; mapping functional regions of the protein; and in studying associated physiological pathways. In addition, modulation of the gene activity <i>in vivo</i> is used for prophylactic and therapeutic purposes.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## NOVEL MAMMALIAN G-PROTEIN COUPLED RECEPTORS HAVING EXTRACELLULAR LEUCINE RICH REPEAT REGIONS

### INTRODUCTION

#### 5 Field of the Invention

The field of this invention is the G-protein coupled receptor family of proteins.

#### Background

Gonadotropins (Luteinizing hormone, LH; follicle stimulating hormone, FSH; chorionic gonadotropin, CG) and thyrotropin (TSH)) are essential for the growth and  
10 differentiation of gonads and thyroid gland, respectively. These glycoprotein hormones bind specific target cell receptors on the plasma membrane to activate the cAMP-protein kinase A pathway.

The receptors for LH, FSH and TSH belong to the large G-protein-coupled, seven-trans-membrane protein family but are unique in having a large N-terminal extra-cellular  
15 (ecto-) domain containing leucine-rich repeats important for interaction with large glycoprotein ligands. Studies suggest that in these receptors, the extra-cellular leucine rich repeat region serves as a "baseball glove" which efficiently catches its corresponding large hormone ligand and optimally orients it for interaction with the seven trans-membrane-helical domain of the receptor.

20 Because hormones and receptors play a prominent role in a variety of physiological processes, there is continued interest in the identification of novel receptors and their ligands, as well as the genes encoding the same.

#### Relevant Literature

References of interest include: el Tayar, N, "Advances in the Molecular  
25 Understanding of Gonadotropins-Receptors Interactions," Mol. Cell. Endocrinol. (December 20, 1996) 125: 65-70; Bhowmick et al., "Determination of Residues Important in Hormone Binding to the Extracellular Domain of the Luteinizing Hormone/Chorionic Gonadotropin Receptor by Site-Directed Mutagenesis and Modeling," Mol. Endocrinol. (September 1996) 10: 1147-1159; Thomas et al., "Mutational Analyses of the  
30 Extracellular Domain of the Full-Length Lutropin/Choriogonadotropin Receptor Suggest Leucine-Rich Repeats 1-6 are Involved in Hormone Binding," Mol. Endocrinol. (June 1996) 10:760-768; Segaloff & Ascoli, "The Gonadotropin Receptors: Insights from the

Cloning of their cDNAs," Oxf. Rev. Reprod. Biol. (1992) 14: 141-168; Braun et al., "Amino-Terminal Leucine-Rich Repeats in Gonadotropin Receptors Determine Hormone Selectivity," EMBO J (July 1991) 10: 1885-1890; and Segaloff et al., "Structure of the Lutropin/Choriogonadotropin Receptor," Recent Prog. Horm. Res. (1990) 46: 261-301.

5

#### SUMMARY OF THE INVENTION

Three novel mammalian G-protein coupled receptors having extra-cellular leucine rich repeat domains, i.e. LGR4, LGR5 and LGR7, and polypeptide compositions related thereto, as well as nucleotide compositions encoding the same, are provided. The subject  
10 proteins, polypeptide and nucleic acid compositions find use in a variety of different applications, including the identification of homologous or related genes; the production of compositions that modulate the expression or function of the subject proteins; in the identification of endogenous ligands for the subject orphan receptors; in the generation of functional binding proteins for the neutralization of the actions of endogenous ligands; in  
15 gene therapy; in mapping functional regions of the protein; and in studying associated physiological pathways. In addition, modulation of the gene activity *in vivo* is used for prophylactic and therapeutic purposes, and the like.

#### BRIEF DESCRIPTION OF THE FIGURES

20 Fig. 1 provides the nucleotide and amino acid sequence for human LGR4.

Fig. 2 provides the nucleotide and amino acid sequence for human LGR5.

Fig. 3 provides the nucleotide and amino acid sequence for human LGR7, long form.

25 Fig. 4 provides the nucleotide and amino acid sequence for human LGR7, short form.

Fig. 5 provides an alignment comparison of the long and short forms of LGR7.

Figs. 6 provides a comparison of deduced amino acid sequence of LGR4 and 5 cDNAs and those encoding FSH and LH receptors.

30

## DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Novel mammalian G-protein coupled receptors having extra-cellular leucine rich repeat regions (i.e. LGR4, LGR5 and LGR7) and polypeptide compositions related thereto, as well as nucleic acid compositions encoding the same, are provided. The subject polypeptide and/or nucleic acid compositions find use in a variety of different applications, including the identification of homologous or related genes; for the identification of endogenous ligands for these novel receptors; the production of compositions that modulate the expression or function of the receptors; for gene therapy; for mapping functional regions of the receptors; in studying associated physiological pathways; for *in vivo* prophylactic and therapeutic purposes; as immunogens for producing antibodies; in screening for biologically active agents; and the like.

Before the subject invention is further described, it is to be understood that the invention is not limited to the particular embodiments of the invention described below, as variations of the particular embodiments may be made and still fall within the scope of the appended claims. It is also to be understood that the terminology employed is for the purpose of describing particular embodiments, and is not intended to be limiting. Instead, the scope of the present invention will be established by the appended claims.

In this specification and the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs.

## CHARACTERIZATION OF LGR4, LGR5 AND LGR7

LGR4, LGR5 and LGR7 are novel mammalian receptors of the G-protein coupled, seven trans-membrane family of proteins, specifically the subfamily of G-protein coupled seven trans-membrane proteins which are characterized by the presence of extra-cellular leucine rich repeat regions. As such, these proteins have trans-membrane segments and extra-cellular regions similar to those found in the known LH, FSH, and TSH receptors. In other words, these proteins have both a G-protein coupled seven trans-membrane region

and a leucine rich repeat extra-cellular domain. The N-terminal extra-cellular domains of these proteins also show high homology with Drosophila Slit and Toll proteins having leucine rich repeats. These proteins are expressed in diverse tissues.

The human LGR4 gene has a nucleotide sequence as shown in SEQ ID NO:01.

- 5 The human LGR4 gene product has an amino acid sequence as shown in SEQ ID NO:02. LGR4 is expressed in a plurality of different tissue types, including ovary, testis, adrenal, placenta, liver, kidney and intestine.

The human LGR5 gene has a nucleotide sequence as shown in SEQ ID NO:03.

- The LGR5 gene product has an amino acid sequence as shown in SEQ ID NO:04. LGR5  
10 has been found to be mainly expressed in muscle, placenta and spinal cord tissue.

- The human LGR7 gene encodes multiple splicing variants, each of which contains a multitude of cysteine-rich low density lipoprotein (LDL) binding motifs at the N-terminus in addition to the leucine rich repeat region. The longer forms of LGR-7 have a higher similarity than shorter forms of LGR-7 to snail LGR in the trans-membrane domain  
15 and the N-terminal LDL binding domain. The overall structure of both the long and short forms of LGR-7 is similar to that of the LH receptor. The human LGR7 short form gene has a nucleotide sequence as shown in SEQ ID NO:05. The LGR7 short form gene product has an amino acid sequence as shown in SEQ ID NO:06. The human LGR7 long form gene has a nucleotide sequence as shown in SEQ ID NO:07. The LGR7 long form  
20 gene product has an amino acid sequence as shown in SEQ ID NO:08. LGR7 is expressed in multiple tissues, including testis, ovary, prostate, intestine and colon.

#### IDENTIFICATION OF *LGR4*, *LGR5* AND *LGR7* SEQUENCES

- Homologs of *LGR4*, *LGR5* and *LGR7* are identified by any of a number of  
25 methods. A fragment of the provided cDNA may be used as a hybridization probe against a cDNA library from the target organism of interest, where low stringency conditions are used. The probe may be a large fragment, or one or more short degenerate primers.

- Nucleic acids having sequence similarity are detected by hybridization under low stringency conditions, for example, at 50°C and 6×SSC (0.9 M sodium chloride/0.09 M  
30 sodium citrate) and remain bound when subjected to washing at 55°C in 1×SSC (0.15 M sodium chloride/0.015 M sodium citrate). Sequence identity may be determined by

hybridization under stringent conditions, for example, at 50°C or higher and 0.1×SSC (15 mM sodium chloride/0.15 mM sodium citrate). Nucleic acids having a region of substantial identity to the provided *LGR4*, *LGR5* and/or *LGR7* sequences, e.g. allelic variants, genetically altered versions of the gene, etc., bind to the provided sequences  
5 under stringent hybridization conditions. By using probes, particularly labeled probes of DNA sequences, one can isolate homologous or related genes. The source of homologous genes may be any species, e.g., primate species, particularly human; rodents, such as rats and mice; canines; felines; bovines; ovines; equines; yeast; nematodes; etc.

Between mammalian species, e.g., human and mouse, homologs have substantial  
10 sequence similarity, e.g. at least 75% sequence identity, usually at least 90%, more usually at least 95% between nucleotide sequences. Sequence similarity is calculated based on a reference sequence, which may be a subset of a larger sequence, such as a conserved motif, coding region, flanking region, etc. A reference sequence will usually be at least about 18 nt long, more usually at least about 30 nt long, and may extend to the complete  
15 sequence that is being compared. Algorithms for sequence analysis are known in the art, such as BLAST, described in Altschul *et al.* (1990), *J. Mol. Biol.* 215:403-10. Unless specified otherwise, all sequence analysis numbers provided herein are as determined with the BLAST program using default settings. The sequences provided herein are essential for recognizing *LGR4*, *LGR5* and *LGR7*-related and homologous proteins in database  
20 searches.

#### *LGR4*, *LGR5* and *LGR7* NUCLEIC ACID COMPOSITIONS

Nucleic acids encoding *LGR4*, *LGR5* and *LGR7* may be cDNA or genomic DNA or a fragment thereof. The terms "*LGR4* gene," "*LGR5* gene" and "*LGR7* gene" shall be  
25 intended to mean the open reading frame encoding specific *LGR4*, *LGR5* and *LGR7* polypeptides, and *LGR4*, *LGR5* and *LGR7* introns, as well as adjacent 5' and 3' non-coding nucleotide sequences involved in the regulation of expression, up to about 20 kb beyond the coding region, but possibly further in either direction. The gene may be introduced into an appropriate vector for extra-chromosomal maintenance or for integration into a  
30 host genome.

The term "cDNA" as used herein is intended to include all nucleic acids that share the arrangement of sequence elements found in native mature mRNA species, where sequence elements are exons and 3' and 5' non-coding regions. Normally mRNA species have contiguous exons, with the intervening introns, when present, removed by nuclear  
5 RNA splicing, to create a continuous open reading frame encoding an LGR4, LGR5 and LGR7 protein.

A genomic sequence of interest comprises the nucleic acid present between the initiation codon and the stop codon, as defined in the listed sequences, including all of the introns that are normally present in a native chromosome. It may further include the 3'  
10 and 5' untranslated regions found in the mature mRNA. It may further include specific transcriptional and translational regulatory sequences, such as promoters, enhancers, *etc.*, including about 1 kb, but possibly more, of flanking genomic DNA at either the 5' or 3' end of the transcribed region. The genomic DNA may be isolated as a fragment of 100 kbp or smaller, and substantially free of flanking chromosomal sequence. The genomic  
15 DNA flanking the coding region, either 3' or 5', or internal regulatory sequences as sometimes found in introns, contains sequences required for proper tissue and stage specific expression.

The sequence of the 5' flanking region may be utilized for promoter elements, including enhancer binding sites, that provide for developmental regulation in tissues  
20 where *LGR4*, *LGR5* and/or *LGR7* is expressed. The tissue specific expression is useful for determining the pattern of expression, and for providing promoters that mimic the native pattern of expression. Naturally occurring polymorphisms in the promoter region are useful for determining natural variations in expression, particularly those that may be associated with disease.

Alternatively, mutations may be introduced into the promoter region to determine  
25 the effect of altering expression in experimentally defined systems. Methods for the identification of specific DNA motifs involved in the binding of transcriptional factors are known in the art, *e.g.* sequence similarity to known binding motifs, gel retardation studies, *etc.* For examples, see Blackwell *et al.* (1995), *Mol. Med.* 1:194-205; Mortlock *et al.*  
30 (1996), *Genome Res.* 6:327-33; and Joulin and Richard-Foy (1995), *Eur. J. Biochem.* 232:620-626.



The regulatory sequences may be used to identify *cis* acting sequences required for transcriptional or translational regulation of *LGR4*, *LGR5* and/or *LGR7* expression, especially in different tissues or stages of development, and to identify *cis* acting sequences and *trans*-acting factors that regulate or mediate *LGR4*, *LGR* and/or *LGR7* expression. Such transcription or translational control regions may be operably linked to an *LGR4*, *LGR5* or *LGR7* gene in order to promote expression of wild type or altered *LGR4*, *LGR5* or *LGR7* or other proteins of interest in cultured cells, or in embryonic, fetal or adult tissues, and for gene therapy.

The nucleic acid compositions of the subject invention may encode all or a part of the subject polypeptides. Double or single stranded fragments may be obtained of the DNA sequence by chemically synthesizing oligonucleotides in accordance with conventional methods, by restriction enzyme digestion, by PCR amplification, *etc.* For the most part, DNA fragments will be of at least 15 nt, usually at least 18 nt or 25 nt, and may be at least about 50 nt. Such small DNA fragments are useful as primers for PCR, hybridization screening probes, *etc.* Larger DNA fragments, *i.e.* greater than 100 nt are useful for production of the encoded polypeptide. For use in amplification reactions, such as PCR, a pair of primers will be used. The exact composition of the primer sequences is not critical to the invention, but for most applications the primers will hybridize to the subject sequence under stringent conditions, as known in the art. It is preferable to choose a pair of primers that will generate an amplification product of at least about 50 nt, preferably at least about 100 nt. Algorithms for the selection of primer sequences are generally known, and are available in commercial software packages. Amplification primers hybridize to complementary strands of DNA, and will prime towards each other.

The *LGR4*, *LGR* and *LGR7* genes are isolated and obtained in substantial purity, generally as other than an intact chromosome. Usually, the DNA will be obtained substantially free of other nucleic acid sequences that do not include an *LGR4*, *LGR5* or *LGR7* sequence or fragment thereof, generally being at least about 50%, usually at least about 90% pure and are typically "recombinant", *i.e.* flanked by one or more nucleotides with which it is not normally associated on a naturally occurring chromosome.

The DNA may also be used to identify expression of the gene in a biological specimen. The manner in which one probes cells for the presence of particular nucleotide

sequences, as genomic DNA or RNA, is well established in the literature and does not require elaboration here. DNA or mRNA is isolated from a cell sample. The mRNA may be amplified by RT-PCR, using reverse transcriptase to form a complementary DNA strand, followed by polymerase chain reaction amplification using primers specific for the subject DNA sequences. Alternatively, the mRNA sample is separated by gel electrophoresis, transferred to a suitable support, *e.g.* nitrocellulose, nylon, *etc.*, and then probed with a fragment of the subject DNA as a probe. Other techniques, such as oligonucleotide ligation assays, *in situ* hybridizations, and hybridization to DNA probes arrayed on a solid chip may also find use. Detection of mRNA hybridizing to the subject sequence is indicative of *LGR4*, *LGR5* and/or *LGR7* gene expression in the sample.

The sequence of an *LGR4*, *LGR5* or *LGR7* gene, including flanking promoter regions and coding regions, may be mutated in various ways known in the art to generate targeted changes in promoter strength, sequence of the encoded protein, *etc.* The DNA sequence or protein product of such a mutation will usually be substantially similar to the sequences provided herein, *i.e.* will differ by at least one nucleotide or amino acid, respectively, and may differ by at least two but not more than about ten nucleotides or amino acids. The sequence changes may be substitutions, insertions, deletions, or a combination thereof. Deletions may further include larger changes, such as deletions of a domain or exon. Other modifications of interest include epitope tagging, *e.g.* with the FLAG system, HA, *etc.* For studies of subcellular localization, fusion proteins with green fluorescent proteins (GFP) may be used.

Techniques for *in vitro* mutagenesis of cloned genes are known. Examples of protocols for site specific mutagenesis may be found in Gustin *et al.* (1993), *Biotechniques* 14:22; Barany (1985), *Gene* 37:111-23; Colicelli *et al.* (1985), *Mol. Gen. Genet.* 199:537-9; and Prentki *et al.* (1984), *Gene* 29:303-13. Methods for site specific mutagenesis can be found in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, CSH Press 1989, pp. 15.3-15.108; Weiner *et al.* (1993), *Gene* 126:35-41; Sayers *et al.* (1992), *Biotechniques* 13:592-6; Jones and Winistorfer (1992), *Biotechniques* 12:528-30; Barton *et al.* (1990), *Nucleic Acids Res* 18:7349-55; Marotti and Tomich (1989), *Gene Anal. Tech.* 6:67-70; and Zhu (1989), *Anal Biochem* 177:120-4. Such mutated genes may

be used to study structure-function relationships of LGR4, LGR5 and/or LGR7, or to alter properties of the protein that affect its function or regulation.

#### LGR4, LGR5 and LGR7 POLYPEPTIDES

- 5        Also provided by the subject invention are LGR4, LGR5 and LGR7 polypeptide compositions. The term polypeptide composition as used herein refers to both the full length proteins as well as portions or fragments thereof. Also included in this term are variations of the naturally occurring proteins, where such variations are homologous or substantially similar to the naturally occurring protein, be the naturally occurring protein
- 10      the human protein, mouse protein, or protein from some other species which naturally expresses an LGR4, LGR5 or LGR7 protein, usually a mammalian species. A candidate homologous protein is substantially similar to an LGR4, LGR5 or LGR7 protein of the subject invention, and therefore is an LGR4, LGR5 or LGR7 protein of the subject invention, if the candidate protein has a sequence that has at least about 80%, usually at
- 15      least about 90% and more usually at least about 98% sequence identity with an LGR4, LGR5 or LGR7 protein, as measured by BLAST, *supra*. In the following description of the subject invention, the term "LGR4, LGR5 or LGR7-protein" is used to refer not only to the human LGR4, LGR5 or LGR7 protein, but also to homologs thereof expressed in non-human species, e.g. murine, rat and other mammalian species.
- 20        The subject gene may be employed for producing all or portions of LGR4, LGR5 and LGR7 polypeptides. By "LGR4 polypeptide/protein", "LGR5 polypeptide/protein," and "LGR7 polypeptide/protein" is meant an amino acid sequence encoded by an open reading frame (ORF) of *LGR4*, *LGR5* and *LGR7* genes, including the full-length native polypeptide and fragments thereof, particularly biologically active fragments and/or
- 25      fragments corresponding to functional domains, e.g. extra-cellular regions; and including fusions of the subject polypeptides to other proteins or parts thereof, e.g. chimeric proteins. For expression, an expression cassette may be employed. The expression vector will provide a transcriptional and translational initiation region, which may be inducible or constitutive, where the coding region is operably linked under the transcriptional control
- 30      of the transcriptional initiation region, and a transcriptional and translational termination

region. These control regions may be native to an *LGR4*, *LGR5* or *LGR7* gene, or may be derived from exogenous sources.

Expression vectors generally have convenient restriction sites located near the promoter sequence to provide for the insertion of nucleic acid sequences encoding  
5 heterologous proteins. A selectable marker operative in the expression host may be present. Expression vectors may be used for the production of fusion proteins, where the exogenous fusion peptide provides additional functionality, i.e. increased protein synthesis, stability, reactivity with defined antisera, an enzyme marker, e.g.  $\beta$ -galactosidase, etc.

10 Expression cassettes may be prepared comprising a transcription initiation region, the gene or fragment thereof, and a transcriptional termination region. Of particular interest is the use of sequences that allow for the expression of functional epitopes or domains, usually at least about 8 amino acids in length, more usually at least about 15 amino acids in length, to about 25 amino acids, and up to the complete open reading frame  
15 of the gene. After introduction of the DNA, the cells containing the construct may be selected by means of a selectable marker, the cells expanded and then used for expression.

LGR4, LGR5 or LGR7 polypeptides may be expressed in prokaryotes or eukaryotes in accordance with conventional ways, depending upon the purpose for expression. For large scale production of the protein, a unicellular organism, such as *E.*  
20 *coli*, *B. subtilis*, *S. cerevisiae*, insect cells in combination with baculovirus vectors, or cells of a higher organism such as vertebrates, particularly mammals, e.g. COS 7 cells, may be used as the expression host cells. In some situations, it is desirable to express the *LGR4*, *LGR5* or *LGR7* gene in eukaryotic cells, where the LGR4, LGR5 or LGR7 protein will benefit from native folding and post-translational modifications. Small peptides can also  
25 be synthesized in the laboratory. Polypeptides that are subsets of the complete LGR4, LGR5 or LGR7 sequence may be used to identify and investigate parts of the protein important for function or to raise antibodies directed against these regions.

For production of the extracellular domain of the LGR4, LGR5 or LGR7 receptor, the anchored receptor approach as described in Osuga et al, Mol. Endocrinol. (1997) 11:  
30 1659-1668 may be employed. Likewise, the chimeric receptor approach described in Kudo et al, J Biol. Chem. (1996) 271; 22470-22478 may be used.

Such peptides find use in the identification of endogenous ligands and in drug screening for agonists and antagonists using methods described in Osuga, supra. Solubilized extracellular domains find use as therapeutic agents, e.g. in the neutralization of the action of endogenous ligands.

5        With the availability of the protein or fragments thereof in large amounts, by employing an expression host, the protein may be isolated and purified in accordance with conventional ways. A lysate may be prepared of the expression host and the lysate purified using HPLC, exclusion chromatography, gel electrophoresis, affinity chromatography, or other purification technique. The purified protein will generally be at  
10    least about 80% pure, preferably at least about 90% pure, and may be up to and including 100% pure. Pure is intended to mean free of other proteins, as well as cellular debris.

The expressed LGR4, LGR5 and LGR7 polypeptides are useful for the production of antibodies, where short fragments provide for antibodies specific for the particular polypeptide, and larger fragments or the entire protein allow for the production of  
15    antibodies over the surface of the polypeptide. Antibodies may be raised to the wild-type or variant forms of LGR4, LGR5 or LGR7. Antibodies may be raised to isolated peptides corresponding to these domains, or to the native protein.

Antibodies are prepared in accordance with conventional ways, where the expressed polypeptide or protein is used as an immunogen, by itself or conjugated to  
20    known immunogenic carriers, e.g. KLH, pre-S HBsAg, other viral or eukaryotic proteins, or the like. Various adjuvants may be employed, with a series of injections, as appropriate. Both polyclonal and monoclonal antibodies may be produced. For monoclonal antibodies, after one or more booster injections, the spleen is isolated, the lymphocytes immortalized by cell fusion, and then screened for high affinity antibody  
25    binding. The immortalized cells, i.e. hybridomas, producing the desired antibodies may then be expanded. For further description, see Monoclonal Antibodies: A Laboratory Manual, Harlow and Lane eds., Cold Spring Harbor Laboratories, Cold Spring Harbor, New York, 1988. If desired, the mRNA encoding the heavy and light chains may be isolated and mutagenized by cloning in *E. coli*, and the heavy and light chains mixed to  
30    further enhance the affinity of the antibody. Alternatives to *in vivo* immunization as a

method of raising antibodies include binding to phage "display" libraries, usually in conjunction with *in vitro* affinity maturation.

#### DIAGNOSTIC USES

5           The subject nucleic acid and/or polypeptide compositions may be used to analyze a patient sample for the presence of polymorphisms associated with a disease state or genetic predisposition to a disease state. Biochemical studies may be performed to determine whether a sequence polymorphism in an *LGR4*, *LGR* or *LGR7* coding region or control regions is associated with disease. Disease associated polymorphisms may include  
10   deletion or truncation of the gene, mutations that alter expression level, that affect the activity of the protein, and the like.

Changes in the promoter or enhancer sequence that may affect expression levels of *LGR4*, *LGR5* or *LGR7* can be compared to expression levels of the normal allele by various methods known in the art. Methods for determining promoter or enhancer  
15   strength include quantitation of the expressed natural protein; insertion of the variant control element into a vector with a reporter gene such as  $\beta$ -galactosidase, luciferase, chloramphenicol acetyltransferase, *etc.*, that provides for convenient quantitation; and the like.

A number of methods are available for analyzing nucleic acids for the presence of  
20   a specific sequence, *e.g.* a disease associated polymorphism. Where large amounts of DNA are available, genomic DNA is used directly. Alternatively, the region of interest is cloned into a suitable vector and grown in sufficient quantity for analysis. Cells that express *LGR4*, *LGR5* or *LGR7* may be used as a source of mRNA, which may be assayed directly or reverse transcribed into cDNA for analysis. The nucleic acid may be amplified  
25   by conventional techniques, such as the polymerase chain reaction (PCR), to provide sufficient amounts for analysis. The use of the polymerase chain reaction is described in Saiki, *et al.* (1985), *Science* 239:487, and a review of techniques may be found in Sambrook, *et al.* Molecular Cloning: A Laboratory Manual, CSH Press 1989, pp.14.2–14.33. Alternatively, various methods are known in the art that utilize oligonucleotide  
30   ligation as a means of detecting polymorphisms, for examples see Riley *et al.* (1990),

*Nucl. Acids Res.* 18:2887-2890; and Delahunty *et al.* (1996), *Am. J. Hum. Genet.* 58:1239-1246.

A detectable label may be included in an amplification reaction. Suitable labels include fluorochromes, *e.g.* fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-X-rhodamine (ROX), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), radioactive labels, *e.g.* <sup>32</sup>P, <sup>35</sup>S, <sup>3</sup>H; *etc.* The label may be a two stage system, where the amplified DNA is conjugated to biotin, haptens, *etc.* having a high affinity binding partner, *e.g.* avidin, specific antibodies, *etc.*, where the binding partner is conjugated to a detectable label. The label may be conjugated to one or both of the primers. Alternatively, the pool of nucleotides used in the amplification is labeled, so as to incorporate the label into the amplification product.

The sample nucleic acid, *e.g.* amplified or cloned fragment, is analyzed by one of a number of methods known in the art. The nucleic acid may be sequenced by dideoxy or other methods, and the sequence of bases compared to a wild-type *LGR4*, *LGR5* or *LGR7* sequence. Hybridization with the variant sequence may also be used to determine its presence, by Southern blots, dot blots, *etc.* The hybridization pattern of a control and variant sequence to an array of oligonucleotide probes immobilized on a solid support, as described in US 5,445,934, or in WO 95/35505 (the disclosures of which are herein incorporated by reference), may also be used as a means of detecting the presence of variant sequences. Single strand conformational polymorphism (SSCP) analysis, denaturing gradient gel electrophoresis (DGGE), and heteroduplex analysis in gel matrices are used to detect conformational changes created by DNA sequence variation as alterations in electrophoretic mobility. Alternatively, where a polymorphism creates or destroys a recognition site for a restriction endonuclease, the sample is digested with that endonuclease, and the products size fractionated to determine whether the fragment was digested. Fractionation is performed by gel or capillary electrophoresis, particularly acrylamide or agarose gels.

Screening for mutations in *LGR4*, *LGR5* or *LGR7* may be based on the functional or antigenic characteristics of the protein. Protein truncation assays are useful in detecting

deletions that may affect the biological activity of the protein. Various immunoassays designed to detect polymorphisms in LGR4, LGR5 or LGR7 proteins may be used in screening. Where many diverse genetic mutations lead to a particular disease phenotype, functional protein assays have proven to be effective screening tools. The activity of the encoded LGR4, LGR5 or LGR7 protein may be determined by comparison with the wild-type protein.

Antibodies specific for LGR4, LGR5 or LGR7 proteins may be used in staining or in immunoassays. Samples, as used herein, include biological fluids such as semen, blood, cerebrospinal fluid, tears, saliva, lymph, dialysis fluid and the like; organ or tissue culture derived fluids; and fluids extracted from physiological tissues. Also included in the term are derivatives and fractions of such fluids. The cells may be dissociated, in the case of solid tissues, or tissue sections may be analyzed. Alternatively a lysate of the cells may be prepared.

Diagnosis may be performed by a number of methods to determine the absence or presence or altered amounts of normal or abnormal LGR4, LGR5 or LGR7 in patient cells. For example, detection may utilize staining of cells or histological sections, performed in accordance with conventional methods. Cells are permeabilized to stain cytoplasmic molecules. The antibodies of interest are added to the cell sample, and incubated for a period of time sufficient to allow binding to the epitope, usually at least about 10 minutes. The antibody may be labeled with radioisotopes, enzymes, fluorescers, chemilumescers, or other labels for direct detection. Alternatively, a second stage antibody or reagent is used to amplify the signal. Such reagents are well known in the art. For example, the primary antibody may be conjugated to biotin, with horseradish peroxidase-conjugated avidin added as a second stage reagent. Alternatively, the secondary antibody conjugated to a fluorescent compound, *e.g.* fluorescein, rhodamine, Texas red, *etc.* Final detection uses a substrate that undergoes a color change in the presence of the peroxidase. The absence or presence of antibody binding may be determined by various methods, including flow cytometry of dissociated cells, microscopy, radiography, scintillation counting, *etc.*

Diagnostic screening may also be performed for polymorphisms that are genetically linked to a disease predisposition, particularly through the use of microsatellite



markers or single nucleotide polymorphisms. Frequently the microsatellite polymorphism itself is not phenotypically expressed, but is linked to sequences that result in a disease predisposition. However, in some cases the microsatellite sequence itself may affect gene expression. Microsatellite linkage analysis may be performed alone, or in combination  
5 with direct detection of polymorphisms, as described above. The use of microsatellite markers for genotyping is well documented. For examples, see Mansfield *et al.* (1994), *Genomics* 24:225-233; Ziegler *et al.* (1992), *Genomics* 14:1026-1031; Dib *et al.*, *supra*.

#### MODULATION OF *LGR4*, *LGR5* and *LGR7* GENE EXPRESSION

10 The *LGR4*, *LGR5* or *LGR7* genes, gene fragments, or the *LGR4*, *LGR5* or *LGR7* protein or protein fragments, are useful in gene therapy to treat disorders associated with *LGR4*, *LGR5* or *LGR7* defects. Expression vectors may be used to introduce the *LGR4*, *LGR5* or *LGR7* gene into a cell. Such vectors generally have convenient restriction sites located near the promoter sequence to provide for the insertion of nucleic acid sequences.  
15 Transcription cassettes may be prepared comprising a transcription initiation region, the target gene or fragment thereof, and a transcriptional termination region. The transcription cassettes may be introduced into a variety of vectors, *e.g.* plasmid; retrovirus, *e.g.* lentivirus; adenovirus; and the like, where the vectors are able to transiently or stably be maintained in the cells, usually for a period of at least about one day, more usually for a  
20 period of at least about several days to several weeks.

The gene or *LGR4*, *LGR5* or *LGR7* protein may be introduced into tissues or host cells by any number of routes, including viral infection, microinjection, or fusion of vesicles. Jet injection may also be used for intramuscular administration, as described by Furth *et al.* (1992), *Anal Biochem* 205:365-368. The DNA may be coated onto gold  
25 microparticles, and delivered intradermally by a particle bombardment device, or "gene gun" as described in the literature (see, for example, Tang *et al.* (1992), *Nature* 356:152-154), where gold microprojectiles are coated with the *LGR4*, *LGR5* or *LGR7* DNA, then bombarded into skin cells.

Antisense molecules can be used to down-regulate expression of *LGR4*, *LGR5*, or  
30 *LGR7* in cells. The anti-sense reagent may be antisense oligonucleotides (ODN), particularly synthetic ODN having chemical modifications from native nucleic acids, or

nucleic acid constructs that express such anti-sense molecules as RNA. The antisense sequence is complementary to the mRNA of the targeted gene, and inhibits expression of the targeted gene products. Antisense molecules inhibit gene expression through various mechanisms, e.g. by reducing the amount of mRNA available for translation, through  
5 activation of RNase H, or steric hindrance. One or a combination of antisense molecules may be administered, where a combination may comprise multiple different sequences.

Antisense molecules may be produced by expression of all or a part of the target gene sequence in an appropriate vector, where the transcriptional initiation is oriented such that an antisense strand is produced as an RNA molecule. Alternatively, the  
10 antisense molecule is a synthetic oligonucleotide. Antisense oligonucleotides will generally be at least about 7, usually at least about 12, more usually at least about 20 nucleotides in length, and not more than about 500, usually not more than about 50, more usually not more than about 35 nucleotides in length, where the length is governed by efficiency of inhibition, specificity, including absence of cross-reactivity, and the like. It  
15 has been found that short oligonucleotides, of from 7 to 8 bases in length, can be strong and selective inhibitors of gene expression (see Wagner *et al.* (1996), *Nature Biotechnol.* 14:840-844).

A specific region or regions of the endogenous sense strand mRNA sequence is chosen to be complemented by the antisense sequence. Selection of a specific sequence  
20 for the oligonucleotide may use an empirical method, where several candidate sequences are assayed for inhibition of expression of the target gene in an *in vitro* or animal model. A combination of sequences may also be used, where several regions of the mRNA sequence are selected for antisense complementation.

Antisense oligonucleotides may be chemically synthesized by methods known in  
25 the art (see Wagner *et al.* (1993), *supra*, and Milligan *et al.*, *supra*.) Preferred oligonucleotides are chemically modified from the native phosphodiester structure, in order to increase their intracellular stability and binding affinity. A number of such modifications have been described in the literature, which alter the chemistry of the backbone, sugars or heterocyclic bases.

30 Among useful changes in the backbone chemistry are phosphorothioates; phosphorodithioates, where both of the non-bridging oxygens are substituted with sulfur;

phosphoroamidites; alkyl phosphotriesters and boranophosphates. Achiral phosphate derivatives include 3'-O'-5'-S-phosphorothioate, 3'-S-5'-O-phosphorothioate, 3'-CH<sub>2</sub>-5'-O-phosphonate and 3'-NH-5'-O-phosphoroamidate. Peptide nucleic acids replace the entire ribose phosphodiester backbone with a peptide linkage. Sugar modifications are also used to enhance stability and affinity. The  $\alpha$ -anomer of deoxyribose may be used, where the base is inverted with respect to the natural  $\beta$ -anomer. The 2'-OH of the ribose sugar may be altered to form 2'-O-methyl or 2'-O-allyl sugars, which provides resistance to degradation without comprising affinity. Modification of the heterocyclic bases must maintain proper base pairing. Some useful substitutions include deoxyuridine for deoxythymidine; 5-methyl-2'-deoxycytidine and 5-bromo-2'-deoxycytidine for deoxycytidine. 5-propynyl-2'-deoxyuridine and 5-propynyl-2'-deoxycytidine have been shown to increase affinity and biological activity when substituted for deoxythymidine and deoxycytidine, respectively.

As an alternative to anti-sense inhibitors, catalytic nucleic acid compounds, *e.g.* ribozymes, anti-sense conjugates, *etc.* may be used to inhibit gene expression. Ribozymes may be synthesized *in vitro* and administered to the patient, or may be encoded on an expression vector, from which the ribozyme is synthesized in the targeted cell (for example, see International patent application WO 9523225, and Beigelman *et al.* (1995), *Nucl. Acids Res.* 23:4434-42). Examples of oligonucleotides with catalytic activity are described in WO 9506764. Conjugates of anti-sense ODN with a metal complex, *e.g.* terpyridylCu(II), capable of mediating mRNA hydrolysis are described in Bashkin *et al.* (1995), *Appl. Biochem. Biotechnol.* 54:43-56.

## GENETICALLY ALTERED CELL OR ANIMAL MODELS FOR LGR4, LGR5 AND LGR7 FUNCTION

The subject nucleic acids can be used to generate transgenic, non-human animals or site specific gene modifications in cell lines. Transgenic animals may be made through homologous recombination, where the normal *LGR4*, *LGR5* or *LGR7* locus is altered. Alternatively, a nucleic acid construct is randomly integrated into the genome. Vectors for stable integration include plasmids, retroviruses and other animal viruses, YACs, and the like.

The modified cells or animals are useful in the study of *LGR4*, *LGR5* and/or *LGR7* function and regulation. For example, a series of small deletions and/or substitutions may be made in the host's native *LGR4*, *LGR5* or *LGR7* gene to determine the role of different exons. Of interest is the use of *LGR4*, *LGR5* or *LGR7* to construct transgenic animal  
5 models for disease states. Specific constructs of interest include anti-sense *LGR4*, *LGR5* or *LGR7*, which will block *LGR4*, *LGR5* or *LGR7* expression, expression of dominant negative *LGR4*, *LGR5* or *LGR7* mutations, and over-expression of *LGR4*, *LGR5* or *LGR7* genes. Where an *LGR4*, *LGR5* or *LGR7* sequence is introduced, the introduced sequence may be either a complete or partial sequence of an *LGR4*, *LGR5* or *LGR7* gene native to  
10 the host, or may be a complete or partial *LGR4*, *LGR5* or *LGR7* sequence that is exogenous to the host animal, *e.g.*, a human *LGR4*, *LGR5* or *LGR7* sequence. A detectable marker, such as *lac Z* may be introduced into the *LGR4*, *LGR5* or *LGR7* locus, where upregulation of *LGR4*, *LGR5* or *LGR7* expression will result in an easily detected change in phenotype.

15 One may also provide for expression of the *LGR4*, *LGR5* or *LGR7* gene or variants thereof in cells or tissues where it is not normally expressed, at levels not normally present in such cells or tissues, or at abnormal times of development. By providing expression of *LGR4*, *LGR5* or *LGR7* protein in cells in which it is not normally produced, one can induce changes in cell behavior, *e.g.* through *LGR4*, *LGR5* or *LGR7* mediated activity.

20 DNA constructs for homologous recombination will comprise at least a portion of the *LGR4*, *LGR5* or *LGR7* gene, which may or may not be native to the species of the host animal, wherein the gene has the desired genetic modification(s), and includes regions of homology to the target locus. DNA constructs for random integration need not include regions of homology to mediate recombination. Conveniently, markers for  
25 positive and negative selection are included. Methods for generating cells having targeted gene modifications through homologous recombination are known in the art. For various techniques for transfecting mammalian cells, see Keown *et al.* (1990), *Meth. Enzymol.* 185:527-537.

For embryonic stem (ES) cells, an ES cell line may be employed, or embryonic  
30 cells may be obtained freshly from a host, *e.g.* mouse, rat, guinea pig, *etc.* Such cells are grown on an appropriate fibroblast-feeder layer or grown in the presence of leukemia

inhibiting factor (LIF). When ES or embryonic cells have been transformed, they may be used to produce transgenic animals. After transformation, the cells are plated onto a feeder layer in an appropriate medium. Cells containing the construct may be detected by employing a selective medium. After sufficient time for colonies to grow, they are picked  
5 and analyzed for the occurrence of homologous recombination or integration of the construct. Those colonies that are positive may then be used for embryo manipulation and blastocyst injection. Blastocysts are obtained from 4 to 6 week old superovulated females. The ES cells are trypsinized, and the modified cells are injected into the blastocoel of the blastocyst. After injection, the blastocysts are returned to each uterine horn of  
10 pseudopregnant females. Females are then allowed to go to term and the resulting offspring screened for the construct. By providing for a different phenotype of the blastocyst and the genetically modified cells, chimeric progeny can be readily detected.

The chimeric animals are screened for the presence of the modified gene and males and females having the modification are mated to produce homozygous progeny. If the  
15 gene alterations cause lethality at some point in development, tissues or organs can be maintained as allogeneic or congenic grafts or transplants, or in *in vitro* culture. The transgenic animals may be any non-human mammal, such as laboratory animals, domestic animals, etc. The transgenic animals may be used in functional studies, drug screening, etc., e.g. to determine the effect of a candidate drug on *LGR4*, *LGR5* or *LGR7* or related  
20 gene activation etc.

#### *IN VITRO* MODELS FOR *LGR4*, *LGR5* or *LGR7* FUNCTION

The availability of a number of components in the G-protein coupled receptor family, as previously described, allows *in vitro* reconstruction of the processes or systems  
25 in which members of this family operate. Two or more of the components, such as the isolated receptor and a potential ligand therefore, may be combined *in vitro*, and the behavior assessed in terms of activation of transcription of specific target sequences; modification of protein components, e.g. proteolytic processing, phosphorylation, methylation, etc.; ability of different protein components to bind to each other. The  
30 components may be modified by sequence deletion, substitution, etc. to determine the functional role of specific domains.

Drug screening may be performed using an *in vitro* model, a genetically altered cell or animal, purified LGR4, LGR5 or LGR7 protein, as well as fragments or portions thereof, e.g. solubilized extra-cellular domain or chimeric receptor proteins comprising the LGR4, LGR5 or LGR7 extra-cellular domain. One can identify ligands or substrates that  
5 bind to and modulate the action of LGR4, LGR5 or LGR7. Areas of investigation include the development of agents that beneficially counter abnormalities related to LGR4, LGR5 or LGR7 and the use of such agents in the therapy.

Drug screening identifies agents that modulate the activity of LGR4, LGR5 or LGR7 function in abnormal cells. Of particular interest are screening assays for agents  
10 that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled *in vitro* protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, and the like. The purified protein may also be used for determination of three-dimensional crystal structure, which can be used for modeling intermolecular interactions, such as GTP binding, *etc.*

15 The term "agent" as used herein describes any molecule, e.g. protein or pharmaceutical, with the capability of altering or mimicking the physiological function of LGR4, LGR5 or LGR7. Generally a plurality of assay mixtures are run in parallel with different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, *i.e.* at  
20 zero concentration or below the level of detection.

In some embodiments, candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly  
25 hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids,  
30 purines, pyrimidines, derivatives, structural analogs or combinations thereof.

Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, *etc.* to produce structural analogs.

Of particular interest in certain embodiments are peptidic agents based on LGR4, LGR5 or LGR7, e.g. solubilized extra-cellular domain or chimeric receptor proteins comprising the LGR4, LGR5 or LGR7 extra-cellular domain, where such agents neutralize the activity of endogenous LGR4, LGR5 or LGR7 ligands, e.g. hormones.

Where the screening assay is a binding assay, one or more of the molecules may be joined to a label, where the label can directly or indirectly provide a detectable signal. Various labels include radioisotopes, fluorescers, chemiluminescers, enzymes, specific binding molecules, particles, e.g. magnetic particles, and the like. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin *etc.* For the specific binding members, the complementary member would normally be labeled with a molecule that provides for detection, in accordance with known procedures.

A variety of other reagents may be included in the screening assay. These include reagents like salts, neutral proteins, e.g. albumin, detergents, *etc.*, that are used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Reagents that improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, *etc.*, may be used. The mixture of components are added in any order that provides for the requisite binding. Incubations are performed at any suitable temperature, typically between 4 and 40°C. Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high-throughput screening. Typically between 0.1 and 1 hours will be sufficient.

Other assays of interest detect agents that mimic LGR4, LGR5 or LGR7 function. For example, an expression construct comprising an *LGR4*, *LGR5* or *LGR7* gene may be introduced into a cell line under conditions that allow expression. The level of LGR4, LGR5 or LGR7 activity is determined by a functional assay, as previously described. In one screening assay, the ability of candidate agents to inhibit or enhance LGR4, LGR5 or LGR7 function is determined. Alternatively, candidate agents are added to a cell that lacks functional LGR4, LGR5 or LGR7, and screened for the ability to reproduce LGR4, LGR5 or LGR7 activity in a functional assay.

The compounds having the desired pharmacological activity may be administered in a physiologically acceptable carrier to a host for treatment, *etc.* The compounds may also be used to enhance *LGR4*, *LGR5* or *LGR7* function. The inhibitory agents may be administered in a variety of ways, orally, topically, parenterally *e.g.* subcutaneously, intraperitoneally, by viral infection, intravascularly, *etc.* Topical treatments are of particular interest. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt.%.  
15

The pharmaceutical compositions can be prepared in various forms, such as granules, tablets, pills, suppositories, capsules, suspensions, salves, lotions and the like. Pharmaceutical grade organic or inorganic carriers and/or diluents suitable for oral and topical use can be used to make up compositions containing the therapeutically-active compounds. Diluents known to the art include aqueous media, vegetable and animal oils and fats. Stabilizing agents, wetting and emulsifying agents, salts for varying the osmotic pressure or buffers for securing an adequate pH value, and skin penetration enhancers can be used as auxiliary agents.

25

#### EXPERIMENTAL

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the subject invention, and are not intended to limit the scope of what is regarded as the invention. Efforts have been made to ensure accuracy with respect to the numbers used (*e.g.* amounts, temperature, concentrations, *etc.*) but some experimental errors and deviations  
30



should be allowed for. Unless otherwise indicated, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees centigrade; and pressure is at or near atmospheric.

5 **Example 1. Identification of LGR4 and LGR5**

Human sequences related to the sea anemone and *Drosophila* glycoprotein hormone receptors were identified from the expression sequence tag database (dbEST) at the National Center for Biotechnology Information by using the BLAST server with the BLOSUM62 protein comparison matrix (Altschul SF *et al*, Nucleic Acids Res (1997)  
10 25:3389-3402). Human ESTs showing high homology to two non-overlapping regions of the gonadotropin receptors were identified. Clones AA312798 and AA298810 were found to encode transmembrane four to five of the putative receptor LGR4 whereas AA460529 and AA424098 encode transmembrane two to three of the putative receptor LGR5. Using these ESTs to further search the GenBank EST division database,  
15 overlapping EST sequences were aligned to obtain the longest open reading frame (ORF) for these receptors.

Based on the longest human ORF, specific primers were designed for PCR amplification of LGR4 and LGR5 cDNA fragments from rat ovary and human placenta, respectively. After hybridization with labeled EST clones and confirmation of DNA  
20 sequences by dideoxy DNA sequencing, specific receptor fragments isolated were used to design primers to prepare sub-cDNA libraries enriched with specific receptor cDNAs. For 5' extension, reverse transcription was performed using rat ovarian and human placenta mRNA preparations and receptor-specific primers. Following second strand synthesis, the enriched cDNA pool was tailed at 5'-ends with specific adaptor sequences to allow further  
25 PCR amplification. For 3' extension, rat ovarian or human placenta mRNAs were reversed transcribed using oligo-dT, followed by second strand synthesis using receptor-specific primers and adaptor tailing. These mini-libraries were further used as templates for PCR amplification of upstream or downstream cDNAs specific for each receptor using internal primers. PCR products with a strong hybridization signal to each receptor cDNA  
30 fragment were subcloned into the pUC18 or pcDNA3 vectors. After screening of these sublibraries based on colony hybridization using specific receptor probes, clones with 5'-

or 3'-sequences of the putative receptors were identified and isolated for DNA sequencing. As needed, the procedure was repeated up to three times to generate cDNAs encoding the complete ORF of each putative receptor for sequence analysis and for the expression of receptor proteins in eukaryotic cells. The entire coding sequences of each gene were also amplified with specific primers flanking the entire ORF in independent experiments. At least three independent PCR clones were sequenced to verify the authenticity of coding sequences. The nucleotide sequence of LGR4, as well as the amino acid sequence of the product encoded by the ORF thereof, is provided in Fig. 1. The nucleotide sequence of LGR5, as well as the amino acid sequence of the product encoded by the ORF thereof, is provided in Fig. 2.

**Example 2. Comparison of deduced amino acid sequence of LGR4 and 5 cDNAs and those encoding FSH and LH receptors.**

Sequence alignment of LGR4 and LGR5 with known human glycoprotein hormone receptors was performed and the results are shown in Fig. 6. Shaded residues are identical in at least two of the four receptor proteins shown.

**Example 3. Expression pattern of LGR4 and 5 mRNA transcripts in different tissues.**

For northern blot analysis, poly (A)<sup>+</sup>-selected RNA from different human tissues was hybridized with a <sup>32</sup>P-labeled cDNA probes. After washing, the blots were exposed to X-ray films at -70C for five days. Subsequent hybridization with a beta-actin cDNA probe was performed to estimate nucleic acid loading (8 h exposure). LGR4 was shown to be expressed in placenta, ovary, testis, adrenal, spinal cord, thyroid, stomach, trachea, heart, pancreas, kidney, prostate and spleen while LGR5 was shown to be expressed in the skeletal muscle, placenta, spinal cord, brain, adrenal, colon, stomach, ovary and bone marrow.

**Example 4. Chromosomal localization of LGR4 and 5 in human.**

Using genomic fragments of LGR4 (>100 Kb) and LGR5 (>100 Kb) as probes, chromosomal localization of these genes were detected using the FISH method to banded DNA in chromosomal 5q34-35.1 and 12q15, respectively.

5

**Example 5. Identification of LGR7.**

Analysis of EST databases has revealed a novel LGR closely related to a G protein-coupled receptor from pond snail (*Lymnaea stagnalis*, accession no. 481946). Because the snail G-protein coupled receptor shared the leucine-rich repeat ectodomain and seven  
10 transmembrane region characteristics of mammalian LGRs, the novel EST sequence could encode either a homologue of snail receptor or a novel mammalian LGR. For the isolation of LGR7 cDNA, a Clontech Marathon-ready testis cDNA pool was used as the template for 5' and 3' RACE with adapter and gene-specific primers. Sequence analysis of the RACE products showed that LGR7 gene encode at least two splicing variants differ at the N-  
15 terminus. The nucleotide sequence of the long variant, as well as the amino acid sequence of the product encoded by the ORF thereof, is provided in Fig. 3; while the nucleotide sequence of the short variant, as well as the amino acid sequence of the ORF thereof, is provided in Fig. 4. Both variants contain a classical C-terminal 7-transmembrane region and a leucine-rich repeat ectodomain flanked by cysteine rich regions found in other mammalian LGRs. The  
20 long form LGR7 contains extra 35 amino acids in the N-terminal cysteine rich region as compared to the short form LGR7. Of interest, analysis of the LGR7 ORF from either variant showed that its tertiary structure resembles that of mammalian LGRs instead of the snail receptor, which shares the greatest identity in the transmembrane region. These findings suggest that LGR7 and snail receptor diverged early during evolution and LGR7 perhaps  
25 adopted new function in higher organisms.

Based on the LGR7 cDNA sequence, we further identified a human genomic DNA fragment (AQ053279) in the genomic survey sequence division of GenBank that contains part of the LGR7 gene. The authenticity of this genomic clone was confirmed by Southern blot hybridization and the genomic clone was used as the probe to identify the  
30 chromosomal localization for LGR7 gene.

It is evident from the above discussion and results that three novel mammalian G-protein coupled receptors, as well as a nucleic acids encoding the same, are provided by the subject invention. The inventions described above find use in a variety of applications, including research and therapeutic applications.

5

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. The publications discussed herein are provided solely for their disclosure prior to the filing date of the  
10 present application. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such a disclosure by virtue of prior invention.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily  
15 apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

## WHAT IS CLAIMED IS:

1. An isolated nucleic acid encoding a mammalian protein selected from the group consisting of LGR4, LGR5 or LGR7.  
5
2. An isolated nucleic acid according to Claim 1, wherein said mammalian protein has the amino acid sequence of SEQ ID NO:2, SEQ ID NO:04, SEQ ID NO:06 or SEQ ID NO:08.
- 10 3. An isolated nucleic acid according to Claim 1, wherein said mammalian protein has an amino acid sequence that is substantially identical to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:04, SEQ ID NO:06 or SEQ ID NO:08.
4. An isolated nucleic acid according to Claim 1, wherein the nucleotide  
15 sequence of said nucleic acid has the sequence selected from the group consisting of: (a) SEQ ID NO:1 or the complementary sequence thereof; (b) SEQ ID NO:03 or the complementary sequence thereof; (c) SEQ ID NO:05 or the complementary sequence thereof; and (d) SEQ ID NO:07 or the complementary sequence thereof.
- 20 5. An isolated nucleic acid comprising at least 18 contiguous nucleotides of the sequence selected from the group consisting of: (a) SEQ ID NO:1 or the complementary sequence thereof; (b) SEQ ID NO:03 or the complementary sequence thereof; (c) SEQ ID NO:05 or the complementary sequence thereof; and (d) SEQ ID  
25 NO:07 or the complementary sequence thereof.
6. An isolated nucleic acid comprising at least 50 contiguous nucleotides of the sequence selected from the group consisting of: (a) SEQ ID NO:1 or the complementary sequence thereof; (b) SEQ ID NO:03 or the complementary sequence thereof; (c) SEQ ID NO:05 or the complementary sequence thereof; and (d) SEQ ID  
30 NO:07 or the complementary sequence thereof.

7. An isolated nucleic acid that hybridizes under stringent conditions to a nucleic acid having the nucleotide sequence selected from the group consisting of: (a) SEQ ID NO:1 or the complementary sequence thereof; (b) SEQ ID NO:03 or the complementary sequence thereof; (c) SEQ ID NO:05 or the complementary sequence thereof; and (d) SEQ ID NO:07 or the complementary sequence thereof.

8. An expression cassette comprising a transcriptional initiation region functional in an expression host, a nucleic acid having a sequence of the isolated nucleic acid according to Claim 1 under the transcriptional regulation of said transcriptional initiation region, and a transcriptional termination region functional in said expression host.

9. A cell comprising an expression cassette according to Claim 8 as part of an extrachromosomal element or integrated into the genome of a host cell as a result of introduction of said expression cassette into said host cell, and the cellular progeny of said host cell.

10. A method for producing a mammalian protein selected from the group consisting of LGR4, LGR5 and LGR7, said method comprising:  
growing a cell according to Claim 9, whereby said mammalian protein is expressed; and  
isolating said protein substantially free of other proteins.

11. A purified polypeptide composition comprising at least 50 weight % of the protein present as a mammalian protein selected from the group consisting of LGR4, LGR5 and LGR7, or a fragment thereof.

12. An antibody binding specifically to a mammalian protein selected from the group consisting of LGR4, LGR5 and LGR7.

13. The antibody of Claim 12, wherein said antibody is a monoclonal antibody.

14. A non-human transgenic animal model for *LGR4*, *LGR5* or *LGR7* gene function, wherein said transgenic animal comprises an introduced alteration in an *LGR4*,  
5 *LGR5* or *LGR7* gene.

15. The animal model of claim 14, wherein said animal is heterozygous for said introduced alteration.

10 16. The animal model of claim 14, wherein said animal is homozygous for said introduced alteration.

17. The animal model of claim 14, wherein said introduced alteration is a knockout of endogenous *LGR4*, *LGR5* or *LGR7* gene expression.

15

18. A method of screening a sample for the presence of a ligand for a receptor selected from the group consisting of *LGR4*, *LGR5* and *LGR7*, said method comprising:  
contacting said sample with a receptor selected from the group consisting of *LGR4*, *LGR5* and *LGR7* or a mimetic thereof, and

20 detecting the presence of a binding event between said receptor and ligand in said sample.

## &gt;LGR4 nucleotide sequence (SEQ ID NO:01)

ATGCCGGGCCCCGCTAGGGCTGCTCTGCTTCCTCGCCCTGGGGCTGCTCGGCTCGGCCGGGCCAGCGGCGGGCGCCGCCCT  
CTCTGCGCGGGCCCCCTGCAGCTGCGACGGCGACCGCTCGGGTGGACTGCTCCGGAAGGGGTTGACGGCCGTACCGGAGGGT  
CTCAGCGCCTTCACCAAGCACTGGATATCAGTATGAACAATATCACCCAGTTACCAGAAGATGCATTTAAGAGTTTCCCA  
TTTCTAGAGGAGCTACAACCTGGCTGGTAACGACCTTTCTCTATCCATCCAAAAGCCTTGCTGGGCTGAAAAGAACTCAA  
GTCCTAACATCCAGAATAATCAGTTGAGAACAGTGCCCACTGAAGCCATTACGGACTGAGTGCTTTGCAGTCTTTACGC  
TTAGATGCCAACCATATTACCTCAGTCCCGGAGGACAGTTTGAAGGGCTTGCCAGTTACGCCATCTGTGGCTGGATGAC  
AACAGCTTGACGGAAGTGCCCGTGGTCCCCTCAGCAACCTGCCAACCTGCAGGCGCTGACCTTGGCTCTCAACAACATC  
TCAAGCATCCCTGACTTCGCTTTCACCAACCTTCAAGCTTGGTGGTCTGCATCTGCATAACAATAAAATTAAGCCCTC  
AGTCAACACTGTTTTGATGGACTAGATAACCTGGAAACCTTGGACTTGAATTACAATTACTTGGATGAGTTTCCTCAGGCT  
ATTAAAGCCCTTCCCAGCCTTAAAGAGCTGGGATTTACAGTAATCTATTTCTGTTATTCCTGATGGAGCATTGTGGTGGT  
AATCCACTGCTAAGAACTATTCATTTGTATGATAATCCTCTGTCTTTGTGGGGAACCTCAGCATTTCACAACTGTCTGAT  
CTGCATTGCTTAGTCATTTCGTGGTGCAAGCCTGGTGCAGTGGTTCCCAATCTGACCGGAACCTGTCCATTTGGAGAGTCTA  
ACCTTGACAGGGACAAAAATAAGCAGCATACCTGATGATCTGTGCCAAAACCAAAGATGCTGAGGACTCTGGACTTATCT  
TATAACAAATATAAGAGACCTTCCAAGTTTTAATGGTTGTCTGTGCATTTGGAAGAAATTTCAATGCAGCGTAATCAAATCTCC  
CTAATAAAGGAAAATACTTTTCAAGGCCTAACATCTCTAAGGATTCTAGATCTGAGTAGAAACCTGATCCGTGAAATTCAC  
ACTGGAGCTTTTGGGAAGCTTGGGACAATTACTAACCTGGATGTAAGTTTCAATGAATTAACCTTCAATTTCTACGGAAGGC  
CTAAATGGGCTCAATCAACTAAAGCTTGTGGGTAACCTCAAGCTGAAAGACGCCCTGGCAGCCAGAGACTTTGCTAATCTC  
AGGTCTCTATCACTACCATATGCTTATCAGTGTTGTGCATTTTGGGGGTGTGACTCTTATGCAAATTAACACAGAAAGAT  
AACAGCCGCGAAGACACAGTGTGACAAAAGAGAAAGGTGCTACAGATGCAGCAAATGTCACCAGCACTGCTGAGAACGAA  
GAACATAGCCAAATAATATCCACTGTACACCTTCAACAGGTGCTTCAAGCCCTGTGAATATTTACTGGGAAGCTGGATG  
ATTCGCTTACAGTGTGGTTCAATTTCTGGTGCCTTGCTTTTCAACCTGCTTGTCAATTTAACAGTGTGTGCGTCTGT  
TCATCACTGCTTGGCTCCAACTCTTCATAGGCTTGATTTCTGTGCTCACTTACTCATGGGCATCTATACTGGCATCTCT  
ACTTTTCTTATGCTGTGTCTTGGGGCCGATTGCGCAATTTGGCATTTGGTGGGAACTGGCAGCGGCTGCAAGGTAGCC  
GGGTCTCTGCGAGTCTTCTCCTCAGAGAGCGCTGTATTCCTATTAACACTGGCAGCTGTGGAAGAAGCGTATTGCAAG  
GATTTGATGAAACACGGGAAGAGCAGTCACCTCAGACAGTTCAGGTGGCCGCCCTCTAGCTTTGCTGGGTGCCGAGTG  
GCAGGCTGCTTCCCTTTTCCACGGAGGGCAATATTCTGCATCGCCCTTGTGTTTGGCGTTTCTACAGGAGAAACCCCA  
TCGTTAGGATTCACTGTGACCTTAGTGCTATTAACTCACTGGCATTTTACTAATGGCCATTATCTACACTAACTATAC  
TGCAACTTAGAGAAGGAGGACCTGTGCGAAAACCTCCAGTCTAGCGTGATTAAGCACGTTGCCTGGCTCATCTTCACAAAC  
TGCATCTTCTTCTGCCCTGTTGCATTTTCTCATTGCAACATTGATCACGGCAATCTCCATCAGCCCCGAGATAATGAAG  
TCTGTTACACTGATATTCTTCCGTTGCCCTGCTTGCTGAATCCGGTCTGTATGTTTCTTCAACCCAAAGTTTAAAGAA  
GACTGGAAGCTACTGAAGCGGCGTGTTACCGGAAACACGGATCTGTTTCAAGTTTCCATCAGCAGCCAAAGCGGTTGTGGG  
GAACAGGATTTCTACTATGACTGTGGCATGTATTCCACTTGCAGGGTAACCTGACTGTCTGTGACTGCTGTGAGTCATTT  
CTTTTGACAAAACAGTATCATGCAAACTTAATAAAATCGCACAGTTGTCCTGTATTGACAGCGGCTCTTGCCAGAGG  
CCAGAGGCTACTGGTCTGATTGTGGTACACAGTCAGCCATTCTGACTATGCAGATGAAGAAGATTCTTTGTCTCAGAC  
AGCTCTGACCGAGGTCCAGGCCTGTGGACGAGCCTGCTTCTACCAGAGTCGTGGATTCCCTCTGGTGCCTATGCTTACAAT  
CTACAGAGAGTCAGAGACTGA

## &gt;LGR4 amino acid sequence (SEQ ID NO:02)

MPGPLGLLCLFALGLLGSAGPSGAAPPLCAAPCSCDGDRRVDCSGKGLTAVPEGLSAFTQALDISMNNITQLPEDAFKSF  
FLEELQLAGNDLSLIHPKALSGLKELKVLTLQNNQLRTPSEAIHGLSALQSLRLDANHITSVPEDSFEGVLQRLHLWDD  
NSLTEVPVRPLSNLPTLQALTALNNISSIPDFAFTNLSSLVVLHLHNNKIKSLSQHCFDGLDNLETLDLNYNYLDEFQA  
IKALPSLKELGFHSNSISVIPDGAFFGNPLLRTIHLNDPLSFVGNFAFHNLSDLHCLVIRGASLVQWFNLTGTVHLESL  
TLTGTKISSIPDDLQCNQKMLRLDLSYNNIRDLPFNGCRALIEISLQRNQISLIKENTFQGLTSLRILDLNRNLIREIH  
SGAFAKLGTITNLDSFNELTSPFTEGLNGLNQLKLVGNFKLDALAARDFANLRSLSVPYAYQCAFWGCDLCKLNTED  
NSPQEHSVTKEKGATDAANVTSTAENEHSQIIHCTPSTGAFKPCYLLGSWMIRLTWVFI FLVALLFNLLVILTVFASC  
SSLPASKLFIGLISVSNLLMGIYTGILTFDAVSWGRFAEFGIWWETGSGCKVAGSLAVFSSESASFLLTLAAVERSVFAK  
DLMKHGKSSHLRQFQVAALLALLGAAVAGCFPLFHGGQYASPLCLPFPTGETPSLGFVTVLVLLNSLAFLLMAIIYTKLY  
CNLEKEDLSENSQSSVIKHWLIFTNCIFFCPVAFFSFAPLITAISISPEIMKSVTLIFFPLPACLPNVLYVFFNPKFKE  
DWKLLKRRVTRKHGSVSVSISQGGCGEQDFYDCGMYSHLQGNLTVDCCESFLLTKPVSCKHLIKSHSCPVLTAASCQR  
PEAYWSDCGTQSAHSDYADEEDSFVSDSSDQVQACGRACFYQSRGFPLVRYAYNLQVRD

FIG. 1



**>Nucleotide sequence of LGR5 (total 2082 nucleotides) (SEQ ID NO:03)**

CTACATCTCCATAACAATAGAATCCACTCCCTGGGAAAGAAATGCTTTGATGGGCTCCACAGCCTAGAGACTTTAGATTTA  
AATTACAATAACCTTGATGAATTCCCCACTGCAATTAGGACACTCTCCAACCTTAAAGGAAGTAGGATTTATAGCAACAAT  
ATCAGGTCGATACCTGAGAAAGCATTGTAGGCAACCCCTTCTCTTATTACAATACATTTCTATGACAATCCCATCCAATTT  
GTTGGGAGATCTGCTTTTCAACATTTACCTGAACTAAGAACTGACTCTGAATGGTGCCCTACAAATAACTGAATTTCT  
GATTTAACTGGAAGTGCAACCTGGAGAGTCTGACTTTAACTGGAGCACAGATCTCATCTCTTCTCAAACCGTCTGCAAT  
CAGTTACCTAATCTCCAAGTGCTAGATCTGTCTTACAACCTATTAGAAGATTTACCCAGTTTTTCAGTCTGCCAAAAGCTT  
CAGAAAATTGACCTAAGACATAATGAAATCTACGAAATTAAGTTGACACTTTCCAGCAGTTGCTTAGCCTCCGATCGCTG  
AATTTGGCTTGGAAACAAATTGCTATTATTACCCCAATGCATTTTCCACTTTGCCATCCCTAATAAGCTGGACCTATCG  
TCCAACCTCTGTCGTCTTTTCTATAACTGGGTTACATGGTTTAACTCACTTAAATTAACAGGAAATCATGCCTTACAG  
AGCTGGATATCATCTGAAACTTTCCAGAACTCAAGGTATAGAAATGCCTTATGCTTACCAGTGCTGTGCATTTGGAGTG  
TGTGAGAATGCCTATAAGATTTCTAATCAATGGAATAAAGGTGACAACAGCAGTATGGACGACCTTCATAAGAAAGATGCT  
GGAATGTTTTCAGGCTCAAGATGAACGTGACCTTGAAGATTTCTGCTTGACTTTGAGGAAGACCTGAAAGCCCTTCATTCA  
GTGCAGTGTTACCTTCCCCAGGCCCTTCAAACCTGTGAACACCTGCTTGATGGCTGGCTGATCAGAATTGGAGTGTGG  
ACCATAGCAGTTCTGGCACTTACTTGTAAATGCTTTGGTGACTTCAACAGTTTTTCAGATCCCTCTGTACATTTCCCCCATT  
AACTGTTAATTGGGGTTCATCGCAGCAGTGAACATGCTCACGGGAGTCTCCAGTGCCGTGCTGGCTGGTGTGGATGCGTTC  
ACTTTTGGCAGCTTTGCACGACATGGTGCTGGTGGGAGAATGGGGTTGGTTGCCATGTCATGGTTTTTTGTCCATTTT  
GCTTCAGAATCATCTGTTTTCTGCTTACTCTGGCAGCCCTGGAGCGTGGGTTCTCTGTGAAATATTCTGCAAAATTGAA  
ACGAAAGCTCCATTTTCTAGCCTGAAAGTAATCATTTTGTCTGTGCCCTGTGCGCTTGACCATGGCCGAGTTCCTCTG  
CTGGGTGGCAGCAAGTATGGCGCCTCCCTCTCTGCTGCTTTGCTTTTGGGGAGCCAGCACCATGGGCTACATGGTC  
GCTCTCATCTTGCTCAATTCCTTTGCTTCTCATGATGACCATTGCCTACACCAAGCTCTACTGCAATTTGGACAAGGGA  
GACCTGGAGAATATTTGGGACTGCTCTATGGTAAACACATTGCCCTGTTGCTCTTACCACACTGCATCCTAAACTGCCCT  
GTGGCTTTCTTGCTCTCTCTCTTTAATAAACCTTACATTTATCAGTCTGAAGTAATTAAGTTTATCCTTCTGGTGGA  
GTCCCACTTCTGCATGTCTCAATCCCTTCTCTACATCTTGTTCATCTCACTTTAAGGAGGATCTGGTGAGCCTGAGA  
AAGCAAACCTACGTCTGGACAAGATCAAACACCCAAGCTTGATGTCAATTAAGTCTGATGATGTCGAAAAACAGTCTGT  
GACTCAACTCAAGCCTTGTAACCTTTACCAGCTCCAGCATCACTTATGACCTGCCTCCAGTTCCGTGCCATCACCAGCT  
TATCCAGTGACTGAGAGCTGCCATCTTCTCTGTGGCATTTGTCCCATGTCTCTAA

**>amino acid sequence of LGR5 (total 693 amino acids) (SEQ ID NO:04)**

LHLHNNRIHSLGKKCFDGLHSLETLDLNNLDEFPTAIRTLNLKELGFHSNNIRSIPEKAFVGNPSLITIHFDNPIQF  
VGRSAFQHLPELRTLTLNGASQITEFPDLTGATANLESLLTGQAQISSLPQTVCNQLPNLQVLDLSYNLLEDLPSPVCQKL  
QKIDLRHNEIYEIKVDTFQQLSLRSLNLAWNKIAIHPNAFSTLPSLIKLDLSSNLLSSFPITGLHGLTHLKLGNHALQ  
SWISSENFPELKVIEMPYAYQCAFVCENAYKISNOWNKGDNSMDDLHKKDAGMFQAQDERDLEDLDFEEDLKALHS  
VQCSPSPGPFKPCHELLDGWLIRIGVWTIAVLALTCNALVTSTVFRSPLYISPIKLLIGVIAAVNMLTGVSSAVLAGVDAF  
TFGSFARHGAWWENGCHVIGFLSIFASESSVFLTLAALERGFVSVKYSKAFETKAPFSSSLKVIILLCALLALTMAAVPL  
LGGSKYGASPLCLPLPFGEPTMGYVALILLNSLCFLMMTIAYTKLYCNLDKGDLENIWDCSMVKHIALLLFTNCILNCP  
VAFLSFSSSLINLTFISPEVIKFIILLVVVPLPACLNPLLYILFNPHFKEDLVSLRKQTVVWTRSKHPSLMSINSDDVEKQSC  
DSTQALVTFSTSSITYDLPPSSVPSPAYPVTESCHLSSVAFVPC

**FIG. 2**

>Final LGR7 (LGR7-Long variant) full length sequence (2467 nt) (SEQ ID NO:05).

GAAAGGAGGAAAGAAAAAAGAGGAATGGAAAGAGACAGAGAAAGGAAATGGGAGTGGAAAGGAGGGAGGACTGCTTT  
 GTAACCTGCTAAGATTGCAGACAGAAATAGCACACAACCACTGTGAGCTGTATGCGATTTCAGAAACCAAGACCAAAT  
 TTGCTCACTTTTCAATTAATCAGTTGCTCAGATAGAAGGAAATGACATCTGGTTCTGTCTTCTACATCTTAATTTT  
 TGGAAATATTTTTCTCATGGGGGTGGACAGGATGTCAAGTGCTCCCTTGGCTATTTCCCTGTGGGAACATCACAA  
 AGTGCTTGCCTCAGCTCCTGCACTGTAACGGTGTGGACGACTGCGGAATCAGGCCGATGAGGACAACCTGTGGAGAC  
 AACATGGATGGTCCATGCAATTTGACAAATATTTTGCCAGTTACTACAAAATGACTTCCCAATATCTTTTGAGGC  
 AGAAACACCTGAATGTTTGGTCGGTTCTGTGCCAGTGCAATGTCTTGCCAAGGTCTGGAGCTTGACTGTGATGAAA  
 CCAATTTACGAGCTGTTCCATCGGTTTCTTCAAATGTGACTGCAATGTCACTTCAGTGGAACCTTAATAAGAAAGCTT  
 CCTCCTGATTGCTTCAAGAATTATCATGATCTTCAGAAGCTGTACCTGCAAAACAATAAGATTACATCCATCTCCAT  
 CTATGCTTTTCAGAGGACTGAATAGCCTTACTAACTGTATCTCAGTCATAACAGAATAACCTTCTGAAGCCGGGTG  
 TTTTTGAAGATCTTCACAGACTAGAATGGCTGATAATTGAAGATAATCACCTCAGTCGAATTTCCCCACCAACATTT  
 TATGGACTAAATTTCTTATTCTTCTAGTCTGATGAATAACGTCCTCACCCTTTTACCTGATAAACCTCTCTGTCA  
 ACACATGCCAAGACTACATTGGCTGGACCTTGAAGGCAACCATATCCATAATTTAAGAAATTTGACTTTTATTTCT  
 GCAGTAATTTAACTGTTTTAGTGATGAGGAAAAACAAATTAATCACTTAAATGAAAATACTTTTGACCTCTCCAG  
 AAATCGGATGAATTGGATTTAGGAAGTAATAAGATTGAAAATCTTCCACCGCTTATATTCAAGGACCTGAAGGAGCT  
 GTCACAATTGAATCTTTCTATAATCCAATCCAGAAAATTCAGCAAACCAATTTGATTATCTTGTCAAACCTCAAGT  
 CTCTCAGCCTAGAAGGGATTGAAATTTCAAATATCCAACAAAGGATGTTTAGACCTCTTATGAATCTCTCTCACATA  
 TATTTTAAGAAATTCAGTACTGTGGGTATGCACCACATGTTTCGAGCTGTAAACCAACACTGATGGAATTTCTATC  
 TCTAGAGAATCTTGGCAAGCATTATTCAGAGAGTATTTGTCTGGGTTGTATCTGCAGTTACCTGCTTTGGAACA  
 TTTTGTCTATTGTCATGCGACCTTATATCAGGTCTGAGAACAGCTGTATGCCATGTCAATCATTTCTCTGTCTGT  
 GCCGACTGCTTAATGGGAATATATTTATCGTGATCGGAGGCTTTGACCTAAAGTTTCGTGGAGAATAACAATAAGCA  
 TCGCAGCTGTGGATGGAGAGTACTCATTTGTGAGGATCTTTGGCCATTCTGTCCACAGAAGTATCAGTTT  
 TACTGTAAACATTTCTGACATTGGAAAAATACATCTGCATTGTCTATCTTTTAGATGTGTGAGACCTGGAAAAATGC  
 AGAACAATTACAGTTCTGATTCTCATTTGGATTACTGTTTTATAGTGGCTTTTATTCCATTGAGCAATAAGGAATT  
 TTTCAAAACTACTATGGCACAATGGAGTATGCTTCCCTCTTATTGAGAGATACAGAAAGTATTGGAGCCCAGA  
 TTTATTGAGTGGCAATTTTCTTGGTATTAATTTGGCCGATTTATCATCATAGTTTTTCTATGGAAGCATGTTT  
 TATAGTGTTCATCAAAGTGCCATAACAGCAACTGAAATACGGAATCAAGTTAAAAAGAGATGATCTTGGCAAACG  
 TTTTCTTTTATAGTATTTTCTGATGCATTATGCTGGATACCCATTTTGTAGTGAAATTTCTTCTACTGCTTCAGG  
 TAGAAATACCAGGTACCATAACCTCTTGGGTAGTGATTTTTATCTGCCATTAAACAGTGCTTTGAACCCAATTCTC  
 TATACTCTGACCACAAGACCATTTAAAGAAATGATTATCGGTTTTTGGTATAACTACAGACAAAGAAAATCTATGGA  
 CAGCAAAGGTGAGAAAACATATGCTCCATCATCTGCGGTGGAAATGTGGCCACTGCAGGAGATGCCACCTGAGT  
 TAATGAAGCCGACCTTTTACATACCCCTGTGAAATGTCACTGATTTCTCAATCAACGAGACTCAATTCCTATTCA  
 TGA

>Final LGR7 (LGR7-long variant, total 757 amino acids)(SEQ ID NO:06)

MTSGSVFFYILIFGKYFSHGGQDVKCSLGYFPCGNITKCLPQLLHCNGVDDCGNQADEDCGDNNGWSMQFDKYFA  
 SYXKMTSQYPFAETPECLVGSVPVQCLCQGLELDCDETNLRAVPSVSSNVTAMSLQWNLRKLPDPCFKNYHDLQK  
 LYLQNNKITSISIIYAFRGLNSLTKLYLSHNRITFLKPGVFEDLHRLLEWLIIEDNHLRSISPPTFYGLNSLILLVLMN  
 NVLTRLPLDKPLCQHMPRLHWDLEGNHINLRLNLTIFISCSNLTVLVMRKNKINHLNENTFAPLQKLDELGLSNKIE  
 NLPPLIFKDLKELSQLNLSYNPIQKIQAQFDYLVKLKLSLSEGLEISNIQRMFRPLMNLSHIYFKKFOYCGYAPH  
 VRSCKPNTDGISSLENLLASIIQRVFWVVSATVCFGNIFVICMRPYIRSENKLYAMSIISLCCADCLMGIIYLFVIG  
 GFDLKRGEYNKHAQLWMESTHCQLVGSILAILSTEVSVLLLTFTLEKYICIVYFPFRCPVRPGKCRITITVLILIWITG  
 FIVAFIPLSNKEFFKNYYGTNGVCFPLHSEDTEISGAQIYSVAIFLGINLAAPFIIIVFSYSGSMFYSVHQSAITATEI  
 RNQVKKEMILAKRFFFIVFTDALCWIPFVVKFLSLQVEIPGTITSWVVFILPINSALNPILYLTITRPFKEMIH  
 RFWYNRQRKSMDSKGQKTYAPSFIIWEMWPLQEMPPELMPKPDLTFTYPEMSLISQSTRLNSYS\*

FIG. 3

## &gt;Final LGR7 (LGR7-Short variant) full length sequence (3584 nt)(SEQ ID NO:07)

CTGCTTTGTAAGTCTAAGATTGCAGACAGAAATAGCACACCACTGTGAGCTGTATGCGATTGAGAAACCAAGA  
 CCAAATTTTGCTCACTTTTATTAAATCAGTTGCTCAGATAGAAGGAAATGACATCTGGTTCTGTCTTCTTACATCT  
 TAATTTTGGAAAATATTTTCTCATGGGGTGGACAGGATGTCAAGTGTCTCCCTTGGCTATTTCCCTGTGGGAAC  
 ATCACAAGTGTCTGCTCAGCTCCTGCACTGTAAACGGTGTGGACGACTGCGGGAATCAGGCCGATGAGGACAACTG  
 TGTGTGGTTTTGTGCCAGTGCATGTCTTTGCCAGGTCTGGAGCTTGACTGGATGAAACATTTACGAGTGTTCCAT  
 CGGTTTCTTCAAATGTGACTGCAATGTCACTTCAGTGGAACTTAATAAGAAAGCTTCCCTCTGATTGCTTCAAGAAT  
 TATCATGATCTTTCAGAAGCTGGACCTGCAAAACAATAAGATTACATCCATCTCCATCTATGCTTTCAGAGGACTGAA  
 TAGCCTTACTAACTGTATCTCAGTCATAACAGAATAACCTTCTGAAGCCGGGTGTTTTTGAAGATCTTTCACAGAC  
 TAGAATGGCTGATAAATGAAGATAATCACCTCAGTCGAATTTCCCAACCAACATTTTATGGACTAAATCTCTTATT  
 CTCTTAGTCTGTATGAATAACGTCTCACCCTTACCTGATAAACCTCTCTGTCAACACATGCCAAGACTACATTG  
 GCTGGACCTTGAAGCAACCATATCATAATTTAAGAAATTTGACTTTTATTTCTGTCAGTAATTTAACTGTTTTAG  
 TGATGAGGAAAAACAAAATTAATCACTTAAATGAAAATACTTTTGACCTCTCCAGAACTGGATGAATTGGATTTA  
 GGAAGTAATAAGATTGAAAATCTTCCACCGCTTATATTCAAGGACCTGAAGGAGCTGTCAATTTGAATCTTTCTTA  
 TAATCCAATCCAGAAAATTCAGCAAAACCAATTTGATTATCTTGTCAAACCTCAAGTCTCTCAGCCTAGAAGGGATTG  
 AAATTTCAAATATCCAACAAAGGATGTTTAGACCTCTTATGAATCTCTCTCACATATATTTAAGAAATTCAGTAC  
 TGTGGGTATGCACCACATGTTTCGACGCTGTAAACCAACACTGATGGAATTTATCTCTAGAGAATCTCTTGGAAG  
 CATTATTCAGAGAGTATTTGTCTGGGTGTATCTGCAGTTACCTGCTTTGGAAACATTTTGTGATTTGCATGCGAC  
 CTTATATCAGGTCTGAGAAACAGCTGTATGCCATGTCAATCATTCTCTCTGCTGTGCCGACTGCTTAATGGGAATA  
 TATTTATTCTGATCGGAGGCTTTGACCTAAAGTTTCTGTGGAGAATACAATAAGCATGCGCAGCTGTGGATGGAGAG  
 TACTCATTGTGCTAGCTGTAGGATCTTTGGCCATTCTGTCCACAGAAGTATCAGTTTTACTGTTTAACATTTCTGACAT  
 TGGAAAAATACATCTGCATTGTCTATCTCTTTTAGATGTGTGAGACCTGGAAAATGCAGAACATTAAGTTCTGATT  
 CTCATTGGATTACTGGTTTTATAGTGGCTTTTCACTTTCATTGAGCAATAAGGAATTTTTCAAAAATACTATGGCAC  
 CAATGGAGTATGCTTCCCTCTTCAATCAGAAGATACAGAAAGTATTGGAGCCAGATTTATTCAGTGGCAATTTTTTC  
 TTGGTATTAATTTGGCCGATTTATCATCATAGTTTTTTCCATGGAAGCATGTTTTATAGTGTTCATCAAGTGCC  
 ATAACAGCAACTGAAATACGGAATCAAGTTAAAAAGAGATGATCCTTGCCAAACGTTTTTCTTTATAGTATTTAC  
 TGATGCATTATGCTGGATACCCATTTTGTAGTGAATTTCTTCACTGCTTCAGGTAGAAATACAGGATACCATAA  
 CCTCTTGGGTAGTGATTTTTATTCTGCCCATTAAACAGTGCTTTGAACCAATCTCTTATACTCTGACCACAAGACCA  
 TTTAAAGAAATGATTCATCGGTTTTTGGTATACTACAGACAAAGAAATCTATGGACAGCAAGGTGAGAAAACATA  
 TGCTCCATCATTCTGCGGTGGAATGTGGCCACTGCAGGAGATGCCACCTGAGTTAATGAAGCCGGACCTTTTCA  
 CATACCCTGTGAAATGTCACTGATTTCTCAATCAACGAGACTCAATTCCTATTCTGACTGACTCTGAAATTCATT  
 TCTTCGAGAGAATACTGTGGGGTGCTTCATGAGGGATTTACTGGTATGAAAATGAATACCACAAAATTAATTTAT  
 AATAATAGCTAAGATAAATATTTTACAAGGACATGAGGAAAAATAAAATGACTAATGCTCTTACAAGGGAAGTAA  
 TTATATCAATAATGTATATATATATTAGTAGACATTTTGCATAAGAAATTAAGAGAAATCTACTTCAGTAAGACTTATT  
 CATTTTTCTAACATGCATTTATTGAGTACCCACTACTATGTGCATAGCATTGCAATATAGTCTGGAAGTAGACAGT  
 GCAGAACCTTTCAATCTGTAGATAGTGTTTAATGACAAAAGACTATACAAAGTCCATCTGCAGTTCCTAGTTTAAAG  
 TAGAGCTTTACCTGTATGTGCATCAGCAAGAAATCATAGGCATTTTAAATAAAGGTTTAAAGTTTTGGAATACTCA  
 GTGTATTTGCATCATAGAAAATGTCTGACTGTTTGCAAAATAATATTCTGTTTTAAGAATCCATCTTACCTCTCTTT  
 AAGTTTCCATACACTTGAGAGCCCAACACAATATTTTACTATAAAAGATGCTTTGCTAGAACTCAAAAACAGCA  
 CTTCTTTTGGCACCTTCTGCCCAGTTTCTCTTTGCTTTAAATGAACATCATCATATGGAATTTGGAATAGGAGAGTA  
 TGAGTACGGCAGAGAAGTGGATCAGAAAACTAGAATGAGGATAAACATTTACATTAGTGGAACTCTGAAATAAA  
 TCCTTGTATTGTGAGTAACTGATTTTCAACAAGGATGCCAAGACAAAAGGCTTTTCAACAAACCGTGTGTTTTA  
 AGAACAGACCTAAGTGGTTTAAATCACCCACTTTAGATGGGTGAATGTTATGGTGTGTGAAATATCTCAGTAAAGCA  
 GTTAAAGGAAAAAGAGCTGGAATGCACTGATTGAGGAATTAATTTGAGGAAGGAAAGGTCTGTATGTACACATTT  
 CACTTTAAGCAGAAAATCTTTCTTCAAGAAATGACTTTACTTTCTCTTGCAGCTGCCAGCAGTGAGATACTAACTT  
 TTTAACTAGTTGTTCTTCTCTAGTCTCTACGTTATTAGNATTTTGTGCTTTTATAATGTGAAACCTTTAAGCAGGAG  
 AAGAAAATGTTTTTCAGATAGTTTCAAATACNCCAAAAATGTTTGCAACACAAAAATACTGGAATCNAACCATAATGC  
 CCTTATTGAATATATAGTTGTATAGNTTGTCTGAAAACCC

## &gt;Final LGR7-S ORF (722 amino acids) (SEQ ID NO:08)

MTSGSVFFYILIFGKYFSGGGQDVKCSLGYFP CGNITKCLPQLLHCNGVDDCGNQADEDCNVVVLQCQMSLPGLEL  
 DWMKPFSTVPSVSSNVTAMSLQWNLRKLPDPCKFNHYDLQKLDLQNNKITSISIYAFRGLNSLTCLYLSHNRITFL  
 KPGVFEDLHRLEWLIIEDNHLRSISPPTFYGLNSLILLVLMNVNLTLPDKPLCQHMPRLHLDLEGNHIIHNLRLNT  
 FISCSNLTVLVMRKNKINHLNENTFAPLQKLELDLGSNKIENLPLIFKDLKELS QLNLSYNPIQKIQANQFDYLV  
 KLKSLSLLEGIEISNIQORMFRPLMNLSHIYFKFQYCGYAPHVRSCKPNTDGISSLENLLASIIQRVFVWVVS AVTC  
 FGNIFVICMRPYIRSENKLYAMSIISLCCADCLMGIYLFVIGGFDLFRGEYNKHAQLWMESTHCQLVGLAILSTE  
 VSVLLLTFLTLLEKYICIVYFPFCVRPGKCRITITVLILIIWITGFIVAFIPLSNKEFFKNYYGTNGVCFPLHSEDTESI  
 GAQIYSVAIFLGINLAAFIIVFSYGSFMFYSVHQSATATEIRNQVKEMILAKRFFFIVFTDALCWIPIFVVKFLS  
 LLQVEIPGTITSWVIFILPINSALNPILYTLTRPFKEMIHRFWYNYRQRKSMDSKGQKTYAPSFIVWEMWPLQEM  
 PPELMKPDLFYPCEMSLISQSTRLNSYS\*

FIG. 4

5/8

&gt;Alignment of LGR7-L with LGR7-S

Query=LGR7-L

Sbjct=LGR7-S

Query: 1 MTSGSVFFYILIFGKYFSHGGGQDVKCSLGYFPCGNITKCLPQLLHCNGVDDCGNQADE 60  
 MTSGSVFFYILIFGKYFSHGGGQDVKCSLGYFPCGNITKCLPQLLHCNGVDDCGNQADE

Sbjct: 1 MTSGSVFFYILIFGKYFSHGGGQDVKCSLGYFPCGNITKCLPQLLHCNGVDDCGNQADE 60

Query: 61 NCGDNNGWSMQFDKYFASYIKMTSQYPFEAETPECLVGSVPVQCLCQ---GLELDCDETN 117  
 NC V V C C GLEL +

Sbjct: 61 NC-----VVVLCQCMSLPGLELDWMKP- 82

Query: 118 LRAVPSVSSNVTAMSLQWNLIRKLPPDCFKNYHDLQKLYLQNNKITSISIIYAFRGLNSLT 177  
 +VPSVSSNVTAMSLQWNLIRKLPPDCFKNYHDLQKL LQNNKITSISIIYAFRGLNSLT

Sbjct: 83 FTSVPSVSSNVTAMSLQWNLIRKLPPDCFKNYHDLQKLDLQNNKITSISIIYAFRGLNSLT 142

Query: 178 KLYLSHNRIITFLKPGVFEDLHRLEWLIIEDNHLRSRISPTFYGLNSLILLVLMNNVLTRL 237  
 KLYLSHNRIITFLKPGVFEDLHRLEWLIIEDNHLRSRISPTFYGLNSLILLVLMNNVLTRL

Sbjct: 143 KLYLSHNRIITFLKPGVFEDLHRLEWLIIEDNHLRSRISPTFYGLNSLILLVLMNNVLTRL 202

Query: 238 PDKPLCQHMPRLHWLDLEGNHIIHNLRLNLTIFISCSNLTVLVMRKNKINHLNENTFAPLQKL 297  
 PDKPLCQHMPRLHWLDLEGNHIIHNLRLNLTIFISCSNLTVLVMRKNKINHLNENTFAPLQKL

Sbjct: 203 PDKPLCQHMPRLHWLDLEGNHIIHNLRLNLTIFISCSNLTVLVMRKNKINHLNENTFAPLQKL 262

Query: 298 DELDLGSNKIENLPPLIFKDLKELSQLNLSYNPIQKIQAQFDYLVKLKSLSLLEGIEISN 357  
 DELDLGSNKIENLPPLIFKDLKELSQLNLSYNPIQKIQAQFDYLVKLKSLSLLEGIEISN

Sbjct: 263 DELDLGSNKIENLPPLIFKDLKELSQLNLSYNPIQKIQAQFDYLVKLKSLSLLEGIEISN 322

Query: 358 IQQRMFRPLMNLSHIYFKKFQYCGYAPHVRSCKPNTDGISSLENLLASIIQRVFVWVSA 417  
 IQQRMFRPLMNLSHIYFKKFQYCGYAPHVRSCKPNTDGISSLENLLASIIQRVFVWVSA

Sbjct: 323 IQQRMFRPLMNLSHIYFKKFQYCGYAPHVRSCKPNTDGISSLENLLASIIQRVFVWVSA 382

Query: 418 VTCFGNIFVICMRPYIRSENKLYAMSIISLCCADCLMGIYLFVIGGFDLKRGEYNKHAQ 477  
 VTCFGNIFVICMRPYIRSENKLYAMSIISLCCADCLMGIYLFVIGGFDLKRGEYNKHAQ

Sbjct: 383 VTCFGNIFVICMRPYIRSENKLYAMSIISLCCADCLMGIYLFVIGGFDLKRGEYNKHAQ 442

Query: 478 LWMESTHCQLVGSAILSTEVSLLLLTFLTLEKYICIVYPPFCVRPGKCRITITVLILIWI 537  
 LWMESTHCQLVGSAILSTEVSLLLLTFLTLEKYICIVYPPFCVRPGKCRITITVLILIWI

Sbjct: 443 LWMESTHCQLVGSAILSTEVSLLLLTFLTLEKYICIVYPPFCVRPGKCRITITVLILIWI 502

Query: 538 TGFIVAFIPLSNKEFFKNYYGTNGVCFPLHSEDTE SIGAQIYSVAIFLGINLAAFIIIVF 597  
 TGFIVAFIPLSNKEFFKNYYGTNGVCFPLHSEDTE SIGAQIYSVAIFLGINLAAFIIIVF

Sbjct: 503 TGFIVAFIPLSNKEFFKNYYGTNGVCFPLHSEDTE SIGAQIYSVAIFLGINLAAFIIIVF 562

Query: 598 SYGSMFYSVHQSAITATEIRNQVKEMILAKRFFFIVFTDALCWIPFVVKFSLSLQVEI 657  
 SYGSMFYSVHQSAITATEIRNQVKEMILAKRFFFIVFTDALCWIPFVVKFSLSLQVEI

Sbjct: 563 SYGSMFYSVHQSAITATEIRNQVKEMILAKRFFFIVFTDALCWIPFVVKFSLSLQVEI 622

Query: 658 PGTITSWVVFILPINSALNPILYTLTTRPFKEMIHRFWYNYRQRKSMDSKGQKTYAPSF 717  
 PGTITSWVVFILPINSALNPILYTLTTRPFKEMIHRFWYNYRQRKSMDSKGQKTYAPSF

Sbjct: 623 PGTITSWVVFILPINSALNPILYTLTTRPFKEMIHRFWYNYRQRKSMDSKGQKTYAPSF 682

Query: 718 IWVEMWPLQEMPELMPDLFTYPCEMSLISQSTRLNSYS 757  
 IWVEMWPLQEMPELMPDLFTYPCEMSLISQSTRLNSYS

Sbjct: 683 IWVEMWPLQEMPELMPDLFTYPCEMSLISQSTRLNSYS 722

FIG. 5

6/8

FIG. 6

LGR4 MPPGLGLLCFLALGLLSAGPSGA  
LGR5 MDTSLRGVLLSLPVLQLATG  
LHR MKQRFSAQLLKLQLLPPLPRA  
FSHR MALLVSLLAFLSLGSG  
TSHR MRPADLLQLVLLDLPRDLGG


LGR4	APPL	AA-P	S	DGDR----	RVD	SGKGLTAVPEGLSAFTQA
LGR5	GSSPRSGVLLRG	P-TH	H	EPDGRMLLRVD		SDLGLSELPSNLSVFTSY
LHR	LREAL	P-EP	N	VPDG--ALR-		PGPTAGLTR
FSHR		HHRI	H	SNRVFL----		QESKVTIETPSDLPRNAIE
TSHR	MG	SSPP	E	HQEED--FRVT		KDIQRIPLPPSTQT

[illegible]


LGR4	RTV-	SE	ING	SA	QS	RLDA	H-	TSV	EDS--	FEGLVQLRH	WLD	S-L-	EV	VR
LGR5	RHV-	TE	LQN	RS	QS	RLDA	H-	SYV	P-SC-	FSLGHSRLH	WLD	A-L-	E	VQ
LHR	RYIE	-G	FIN	PQ	KY	SIC-	TG	RKF	DVTKVFSSES	NFI-	EIC	LHI-	T	GN
FSHR	LYIN	-E	FQN	PN	QY	LIS-	TG	KHL	DVHK-IHSLQKVL-	DIQ	INI-	H-	ERN	
TSHR	TYID	-D	LKE	PL	KF	GIF-	TGLKMF	DLTK-VYSTD	IFFI	BIT	PYM-	S	VN	

LGR4 PLSN P-TLQA T AL NISSIPDF T LSS VV H HN K-IKSLSQHC D LDN-LE  
 LGR5 A RS S-ALQAMT AL KIHHPDY G LSSWV H HN R-IHSLGKKC D LHS-LE  
 LHR A QGMNNESTV K YG GFEEVQSH - GTT TS E KE VHLEKMNGA R A-TGPK  
 FSHR S VG SFESVI W NK GIQEIHC - GTQ DE N SD NNLEELPNDV H A-SGPV  
 TSHR A OG CNETLT K YN GFTSVQGY - GTK DAVY NK KYLTVIDKDA G VYSGPS

		←-----→ ←-----→ ←-----→ ←-----→									
LGR4	T	LNYNYLDEF	Q-AIKA	PS	KELGFHSNSISVI	D-GA	GGNPL	RTIH	-	DNPLS	
LGR5	T	LNYYNLDEF	T-AIRT	SN	KELGFHSNNIRSI	E-KA	VGNPS	ITIHF	-	DNPIQ	
LHR	T	ISSTKLQAL	SYGLESIQ	I-ATS-SYSLKKL	SRET	V-N--	LEAT	T	----		
FSHR	I	ISRTRIHSL	SYGLEN	KK	R-ARSTYN-LKKL	TLEKLVA---	MEAS	T	----		
TSHR	L	VSQTSVTAL	SKGLEH	KE	I-ARNTWT-LKKL	LSLS	LH---	TRAD	S	----	



LGR4 FVGNSAFHNLSDLHCLVIRGASLVQWFPNLTGTVHLES~~LT~~LTGTGISSIPDDLCQNQKML  
 LGR5 FVGRSAFQHLPELRTLTNLGASQITEFPDLTG~~TAN~~LES~~LT~~LTGAQISSLPQTVCNQLPNL  
 LHR -----  
 FSHR -----  
 TSHR -----



## C-flank cysteine-rich sequence

LGR4	LVGNFKLKDALAARDFANLRSLSV YAYQ	WGCDSLCKLNTEDNSPQEHSVTKEKGA
LGR5	LTGNHALQSLISSENFPELVKVIEM YAYQ	GVCENAYKISNQWNKGDNSMDDLHKK
LHR	----- --SH	RNLPTKEQNFSSHISENFSKQCESTVR
FSHR	----- --SH	ANWRRQISELHPICNKSILRQEVDMYT
TSHR	----- --SH	KNQKKIRGILESLMCNESSMQSLRQRK

LGR4	TDAANVTSTAENE HS-----
LGR5	DAGMFQAQDERDL DF-----
LHR	KVSNKTLYSSMLA SE-----
FSHR	QTRGQRSSLAEDN SS-----
TSHR	SVNALNSPLHQEY ENLGDSIVGYKEKSKFQDTHNNAHYVVFEEQEDEIIGFGQELKNP

LGR4	-----QIIIH T STGA K	YLLGSWMI
LGR5	-----LLDFEEDLKALHSVQ S SPGP K	HLLDGWLI
LHR	-----LSGWDYIEYGFCPLPKTPR- A EPDA N	DIMGYDFL
FSHR	YSRGFDMTYTEFDYDLCEVVDVT S KPDA N	DIMGYNIL
TSHR	QEETLQAFDSHYDYTCGDSSEDMV T KSDE N	DIMGYKFL

## Transmembrane

	TM 1	TM 2
LGR4	LTV F FLV LLF LL ILTVFA CSS PASKLFIGLISVSNLLM IYTGILTFL AVSW	
LGR5	IGV T AV LTC AL TSTVFR PLYISPIKL IGVIAAVNMLT VSSAVL G AF F	
LHR	VLI L NI IMG MT LFVLLT RYK TVPRF MCNLSFADFCM LYLLLI S SQ K	
FSHR	VLI F SI ITG II LVILTT QYK TVPRF MCNLAFAADLCI IYLLLI S IH K	
TSHR	IVV FVSL LLG VF LLILLT HYK NVPRF MCNLAFAADFCM MYLLLI S LY H	

	TM 3
LGR4	GRFAEFG W E S KV SLA S SA FL LAAV SVFAKDLMKHGKSSH QF
LGR5	GSFARHGAW EN V HVI LSI S FL LAA GFSVKYSAKFET APFSSL
LHR	GQYYNHA D Q S ST FT L YT VIT WHTITYAIHLDQ LR HA
FSHR	SQYHNYA D Q A DA FT L YT AIT WHTITHAMQLDC VQ HA
TSHR	SEYYNHA D Q P NT FT L YT VIT WYAITFAMRLDR IR HA

FIG. 6 (CONT)

8/8

	TM 4			TM 5		
LGR4	QVAALLALLGA	AVAGCF	FHGGQ SASPL	FPTGETPSL	GFTVTLVL	SL LLMA
LGR5	KVIILLCALL	ALTM AV	L G K GASPL	LPFGEPSTM	G MVALIL	SLC LMMT
LHR	ILIMLGGW	LFSSLI ML	V V N MKVSI	F MDVETTL	SQV ILTILI	VV FIIC
FSHR	ASVMVMGW	IFAFAA LF	IF I S MKVSI	MDIDSPL	SQV VMSLLV	VL VVIC
TSHR	CAIMVGGW	VCCFLL LL	V I S AKVSI	MDTETPL	ALA IVFVLT	IV VIVC

	TM 6						
LGR4	II T L	CNL-EKED	SENSQSSVI	HV W	NCIFFC	VA FSFAPLIT	TAIS SPEI
LGR5	IA T L	CNL-DKGD	LENIW CSMV	HI L L	NCILNC	VA LSF SLIN	TLTF SPEV
LHR	AC I I	FAVRNP	ELMATNK	TKIA KM I	DFTCMA	IS FAI AAFK	VPL TVTN
FSHR	GC IHI	LTVRNP	NIVSSSS	TRIA RM M	DFLCMA	IS FAI ASLK	VPL TVSK
TSHR	CCHV I	ITVRNP	QYNPGDK	TKIA RM V	DFICMA	IS YAL AILN	KPL TVSN

	TM 7			
LGR4	M SVTLI	F LPA L	V VF N	
LGR5	I FI LVVV	LPA L	L IL N	
LHR	S VL VL	Y INS A	F AI T	
FSHR	A IL VL	H INS A	F AI T	
TSHR	S IL VL	Y LNS A	F AI T	

**C-terminal tail**

LGR4	PK KE WKL KRRVTRKHG	SVSVSISSQGGCGEQDFYDCGMYSHLQGNLTVCDCCESFL
LGR5	PH KE LVS RKQTYVWTRSKHPSLMSINSDDVEKQSCDSTQALVTFTSSSITYDLPPSS	
LHR	KT QR FFL LSKFGCKRRAELYYRRKDFSAYTSNCKNGFTGSNKPSQSTLKLSTLHCQG	
FSHR	KN RR FFI LSKCGCYEMQAQIYRTETSSSTVHNTHPRNGHCSSAPRVNNGSTYILVPLS	
TSHR	KA QR VFI LSKFGICKRQAQAYRGQRVPPKNSTDIVQVKVTHDMRQGLHNMEDVYELI	

LGR4	LTKPVSCCKHLIKSHSCPVLTAASCQRPEAYWSDCGTQSAHSDYADEEDSFVSDSSDQVQA
LGR5	VPSPAYPVTESCHLSSVAFVPCL
LHR	TALLDKTRYTEC
FSHR	HLAQN
TSHR	ENSHLTPKKQGQISEEYMQTVL

LGR4	CGRACFYQSRGFPLVRYAYNLQVRD
------	---------------------------

**FIG. 6 (CONT)**

## SEQUENCE LISTING

<110> Hsueh, Aaron  
 Hsu, Yu Sheau  
 Liang, Shan-Guang  
 van der Spek, Petrus Johannes

<120> Novel Mammalian G-Protein Coupled  
 Receptors Having Extracellular Leucine Rich Repeat Regions

<130> SUN-84PCT

<160> 8

<170> FastSEQ for Windows Version 3.0

<210> 1  
 <211> 2856  
 <212> DNA  
 <213> human

<400> 1

atgccggggcc	cgctagggct	gctctgcttc	ctcgccctgg	ggctgctcgg	ctcggccggg	60
cccagcggcg	cggcgccgcc	tctctgcgcg	gcgccctgca	gctgcgacgg	cgaccgtcgg	120
gtggactgct	ccggaaaggg	gttgacggcc	gtaccggagg	gtctcagcgc	cttcacccaa	180
gcactggata	tcagtatgaa	caatatcacc	cagttaccag	aagatgcatt	taagagtttc	240
ccattttctag	aggagctaca	actggctggt	aacgaccttt	ctcttatcca	tccaaaagcc	300
ttgtctgggc	tgaaagaact	caaagtccta	acactccaga	ataatcagtt	gagaacagtg	360
cccagtgaag	ccatttcacgg	actgagtgt	ttgcagtcct	tacgcttaga	tgccaaccat	420
attacctcag	tcccggagga	cagttttgaa	gggcttgctc	agttacgcca	tctgtggctg	480
gatgacaaca	gcttgacgga	agtgcctgtg	cgtcccctca	gcaacctgcc	aacctgcag	540
gcgctgacct	tggctctcaa	caacatctca	agcatccctg	acttcgcttt	caccaacctt	600
tcaagcttgg	tggttctgca	tctgcataac	aataaaaatta	aaagcctcag	tcaacactgt	660
tttgatggac	tagataacct	ggaaaacctg	gacttgaatt	acaattactt	ggatgagttt	720
cctcaggcta	ttaaagccct	tcccagcctt	aaagagctgg	gatttcacag	taattctatt	780
tctgttattc	ctgatggagc	atttggtggt	aatccactgc	taagaactat	tcatttgtat	840
gataatcctc	tgtcttttgt	ggggaactca	gcatttcaca	acctgtctga	tctgcattgc	900
ttagtcattc	gtggtgcaag	cctgggtcag	tggttcccca	atctgaccgg	aactgtccat	960
ttggagagtc	taaccttgac	agggacaaaa	ataagcagca	tacctgatga	tctgtgccaa	1020
aaccaaaaga	tgctgaggac	tctggactta	tcttataaca	atataagaga	ccttccaagt	1080
tttaatggtt	gtcgtgcatt	ggaagaaatt	tcattgcagc	gtaatcaaat	ctccctaata	1140
aaggaaaata	cttttcaagg	cctaacatct	ctaaggattc	tagatctgag	tagaaaacctg	1200
atccgtgaaa	ttcacagtgg	agcttttgcg	aaagcttggg	caattactaa	cctggatgta	1260
agttttcaatg	aattaacttc	atttcctacg	gaaggcctaa	atgggctcaa	tcaactaaag	1320
cttggtggta	acttcaagct	gaaagacgcc	ttggcagcca	gagactttgc	taatctcagg	1380
tctctatcag	taccatatgc	ttatcagtgt	tgtgcatttt	gggggtgtga	ctctttatgc	1440
aaattaaaca	cagaagataa	cagcccccaa	gaacacagtg	tgacaaaaga	gaaaggtgct	1500
acagatgcag	caaatgtcac	cagcactgct	gagaacgaag	aacatagcca	aataattatc	1560
cactgtacac	cttcaacagg	tgctttcaag	ccctgtgaat	atttactggg	aagctggatg	1620
attcgcctta	cagtgtggtt	cattttcctg	gtcgccttgc	ttttcaacct	gcttgtcatt	1680
ttaacagtgt	ttgcgtcttg	ttcatcactg	cctgcctcca	aactcttcat	aggcttgatt	1740
tctgtgtcta	acttactcat	gggcatctat	actggcatcc	ttacttttct	tgatgctgtg	1800
tcctggggcc	gatttgccga	atttggcatt	tgggtggaaa	ctggcagcgg	ctgcaaggta	1860
gccgggtctc	tggcagtcct	ctcctcagag	agcgtgtgat	tcctattaac	actggcagct	1920
gtggaaaagaa	gcgtatttgc	aaaggatttg	atgaaacacg	ggaagagcag	tcacctcaga	1980
cagttccagg	tggccgccct	cttagctttg	ctgggtgccg	cagtggcagg	ctgcttcccc	2040
cttttccacg	gagggcaata	ttctgcatcg	cccttggtgt	tgccgtttcc	tacaggagaa	2100



```

accccatcgt taggattcac tgtgacctta gtgctattaa actcactggc atttttacta 2160
atggccatta tctacactaa actatactgc aacttagaga aggaggacct gtcggaaaac 2220
tcccagtcga gcgtgattaa gcacgttgcc tggtcatctc tcacaaactg catcttcttc 2280
tgccctgttg catttttctc atttgacca ttgatcacgg caatctccat cagccccgag 2340
ataatgaagt ctgttacact gatattcttc ccgttgccctg ctgacctgaa tccggctcctg 2400
tatgttttct tcaacccaaa gtttaaagaa gactggaagc tactgaagcg gcgtgttacc 2460
aggaacacag gatctgtttc agtttccatc agcagccaag gcggttggtg ggaacaggat 2520
ttctactatg actgtggcat gtattccac ttgcagggtg acctgactgt ctgtgactgc 2580
tgtgagtcac ttcttttgac aaaaccagta tcatgcaaac acttaataaa atcgcacagt 2640
tgtcctgtat tgacagcggc ctcttgccag aggccagagg cctactggtc tgattgtggt 2700
acacagtcag cccattctga ctatgcagat gaagaagatt cctttgtctc agacagctct 2760
gaccaggtgc aggcctgtgg acgagcctgc ttctaccaga gtcgtggatt ccctctggtg 2820
cgctatgctt acaatctaca gagagtcaga gactga 2856

```

<210> 2  
 <211> 951  
 <212> PRT  
 <213> human

<400> 2

```

Met Pro Gly Pro Leu Gly Leu Leu Cys Phe Leu Ala Leu Gly Leu Leu
 1          5          10          15
Gly Ser Ala Gly Pro Ser Gly Ala Ala Pro Pro Leu Cys Ala Ala Pro
 20          25          30
Cys Ser Cys Asp Gly Asp Arg Arg Val Asp Cys Ser Gly Lys Gly Leu
 35          40          45
Thr Ala Val Pro Glu Gly Leu Ser Ala Phe Thr Gln Ala Leu Asp Ile
 50          55          60
Ser Met Asn Asn Ile Thr Gln Leu Pro Glu Asp Ala Phe Lys Ser Phe
 65          70          75          80
Pro Phe Leu Glu Glu Leu Gln Leu Ala Gly Asn Asp Leu Ser Leu Ile
 85          90          95
His Pro Lys Ala Leu Ser Gly Leu Lys Glu Leu Lys Val Leu Thr Leu
 100          105          110
Gln Asn Asn Gln Leu Arg Thr Val Pro Ser Glu Ala Ile His Gly Leu
 115          120          125
Ser Ala Leu Gln Ser Leu Arg Leu Asp Ala Asn His Ile Thr Ser Val
 130          135          140
Pro Glu Asp Ser Phe Glu Gly Leu Val Gln Leu Arg His Leu Trp Leu
 145          150          155          160
Asp Asp Asn Ser Leu Thr Glu Val Pro Val Arg Pro Leu Ser Asn Leu
 165          170          175
Pro Thr Leu Gln Ala Leu Thr Leu Ala Leu Asn Asn Ile Ser Ser Ile
 180          185          190
Pro Asp Phe Ala Phe Thr Asn Leu Ser Ser Leu Val Val Leu His Leu
 195          200          205
His Asn Asn Lys Ile Lys Ser Leu Ser Gln His Cys Phe Asp Gly Leu
 210          215          220
Asp Asn Leu Glu Thr Leu Asp Leu Asn Tyr Asn Tyr Leu Asp Glu Phe
 225          230          235          240
Pro Gln Ala Ile Lys Ala Leu Pro Ser Leu Lys Glu Leu Gly Phe His
 245          250          255
Ser Asn Ser Ile Ser Val Ile Pro Asp Gly Ala Phe Gly Gly Asn Pro
 260          265          270
Leu Leu Arg Thr Ile His Leu Tyr Asp Asn Pro Leu Ser Phe Val Gly
 275          280          285
Asn Ser Ala Phe His Asn Leu Ser Asp Leu His Cys Leu Val Ile Arg
 290          295          300

```

Gly Ala Ser Leu Val Gln Trp Phe Pro Asn Leu Thr Gly Thr Val His  
 305 310 315 320  
 Leu Glu Ser Leu Thr Leu Thr Gly Thr Lys Ile Ser Ser Ile Pro Asp  
 325 330 335  
 Asp Leu Cys Gln Asn Gln Lys Met Leu Arg Thr Leu Asp Leu Ser Tyr  
 340 345 350  
 Asn Asn Ile Arg Asp Leu Pro Ser Phe Asn Gly Cys Arg Ala Leu Glu  
 355 360 365  
 Glu Ile Ser Leu Gln Arg Asn Gln Ile Ser Leu Ile Lys Glu Asn Thr  
 370 375 380  
 Phe Gln Gly Leu Thr Ser Leu Arg Ile Leu Asp Leu Ser Arg Asn Leu  
 385 390 395 400  
 Ile Arg Glu Ile His Ser Gly Ala Phe Ala Lys Leu Gly Thr Ile Thr  
 405 410 415  
 Asn Leu Asp Val Ser Phe Asn Glu Leu Thr Ser Phe Pro Thr Glu Gly  
 420 425 430  
 Leu Asn Gly Leu Asn Gln Leu Lys Leu Val Gly Asn Phe Lys Leu Lys  
 435 440 445  
 Asp Ala Leu Ala Ala Arg Asp Phe Ala Asn Leu Arg Ser Leu Ser Val  
 450 455 460  
 Pro Tyr Ala Tyr Gln Cys Cys Ala Phe Trp Gly Cys Asp Ser Leu Cys  
 465 470 475 480  
 Lys Leu Asn Thr Glu Asp Asn Ser Pro Gln Glu His Ser Val Thr Lys  
 485 490 495  
 Glu Lys Gly Ala Thr Asp Ala Ala Asn Val Thr Ser Thr Ala Glu Asn  
 500 505 510  
 Glu Glu His Ser Gln Ile Ile Ile His Cys Thr Pro Ser Thr Gly Ala  
 515 520 525  
 Phe Lys Pro Cys Glu Tyr Leu Leu Gly Ser Trp Met Ile Arg Leu Thr  
 530 535 540  
 Val Trp Phe Ile Phe Leu Val Ala Leu Leu Phe Asn Leu Leu Val Ile  
 545 550 555 560  
 Leu Thr Val Phe Ala Ser Cys Ser Ser Leu Pro Ala Ser Lys Leu Phe  
 565 570 575  
 Ile Gly Leu Ile Ser Val Ser Asn Leu Leu Met Gly Ile Tyr Thr Gly  
 580 585 590  
 Ile Leu Thr Phe Leu Asp Ala Val Ser Trp Gly Arg Phe Ala Glu Phe  
 595 600 605  
 Gly Ile Trp Trp Glu Thr Gly Ser Gly Cys Lys Val Ala Gly Ser Leu  
 610 615 620  
 Ala Val Phe Ser Ser Glu Ser Ala Val Phe Leu Leu Thr Leu Ala Ala  
 625 630 635 640  
 Val Glu Arg Ser Val Phe Ala Lys Asp Leu Met Lys His Gly Lys Ser  
 645 650 655  
 Ser His Leu Arg Gln Phe Gln Val Ala Ala Leu Leu Ala Leu Leu Gly  
 660 665 670  
 Ala Ala Val Ala Gly Cys Phe Pro Leu Phe His Gly Gly Gln Tyr Ser  
 675 680 685  
 Ala Ser Pro Leu Cys Leu Pro Phe Pro Thr Gly Glu Thr Pro Ser Leu  
 690 695 700  
 Gly Phe Thr Val Thr Leu Val Leu Leu Asn Ser Leu Ala Phe Leu Leu  
 705 710 715 720  
 Met Ala Ile Ile Tyr Thr Lys Leu Tyr Cys Asn Leu Glu Lys Glu Asp  
 725 730 735  
 Leu Ser Glu Asn Ser Gln Ser Ser Val Ile Lys His Val Ala Trp Leu  
 740 745 750  
 Ile Phe Thr Asn Cys Ile Phe Phe Cys Pro Val Ala Phe Phe Ser Phe  
 755 760 765

Ala Pro Leu Ile Thr Ala Ile Ser Ile Ser Pro Glu Ile Met Lys Ser  
 770 775 780  
 Val Thr Leu Ile Phe Phe Pro Leu Pro Ala Cys Leu Asn Pro Val Leu  
 785 790 795 800  
 Tyr Val Phe Phe Asn Pro Lys Phe Lys Glu Asp Trp Lys Leu Leu Lys  
 805 810 815  
 Arg Arg Val Thr Arg Lys His Gly Ser Val Ser Val Ser Ile Ser Ser  
 820 825 830  
 Gln Gly Gly Cys Gly Glu Gln Asp Phe Tyr Tyr Asp Cys Gly Met Tyr  
 835 840 845  
 Ser His Leu Gln Gly Asn Leu Thr Val Cys Asp Cys Cys Glu Ser Phe  
 850 855 860  
 Leu Leu Thr Lys Pro Val Ser Cys Lys His Leu Ile Lys Ser His Ser  
 865 870 875 880  
 Cys Pro Val Leu Thr Ala Ala Ser Cys Gln Arg Pro Glu Ala Tyr Trp  
 885 890 895  
 Ser Asp Cys Gly Thr Gln Ser Ala His Ser Asp Tyr Ala Asp Glu Glu  
 900 905 910  
 Asp Ser Phe Val Ser Asp Ser Ser Asp Gln Val Gln Ala Cys Gly Arg  
 915 920 925  
 Ala Cys Phe Tyr Gln Ser Arg Gly Phe Pro Leu Val Arg Tyr Ala Tyr  
 930 935 940  
 Asn Leu Gln Arg Val Arg Asp  
 945 950

<210> 3  
 <211> 2082  
 <212> DNA  
 <213> human

<400> 3  
 ctacatctcc ataacaatag aatccactcc ctgggaaaga aatgctttga tgggctccac 60  
 agcctagaga ctttagattt aaattacaat aaccttgatg aattccccac tgcaattagg 120  
 acactctcca acttaaagga actaggattt catagcaaca atatcaggtc gatacctgag 180  
 aaagcatttg taggcaaccc ttctcttatt acaatacatt tctatgacaa tcccatccaa 240  
 ttgtgtggga gatctgcttt tcaacattta cctgaactaa gaacactgac tctgaatggg 300  
 gcctcacaaa taactgaatt tcctgattta actggaactg caaacctgga gagtctgact 360  
 ttaactggag cacagatctc atctcttcct caaacctgct gcaatcagtt acctaacttc 420  
 caagtgcctag atctgtctta caacctatta gaagatttac ccagtttttc agtctgccaa 480  
 aagcttcaga aaattgacct aagacataat gaaatctacg aaattaaagt tgacactttc 540  
 cagcagttgc ttagcctccg atcgtggaat ttggcttgga acaaaattgc tattattcac 600  
 cccaatgcat ttccacttt gccatcccta ataaagctgg acctatcgtc caacctcctg 660  
 tcgtcttttc ctataactgg gttacatggt ttaactcact taaaattaac aggaaatcat 720  
 gccttacaga gctggatc atctgaaaac ttccagaaac tcaaggtnat agaaatgcct 780  
 tatgcttacc agtgctgtgc atttgagtg tgtgagaatg cctataagat ttctaataca 840  
 tggataaaag gtgacaacag cagtatggac gaccttcata agaaagatgc tggaaatggt 900  
 caggctcaag atgaacgtga ccttgaagat ttctgtcttg actttgagga agacctgaaa 960  
 gcccttcatt cagtgcagtg ttacacttcc ccaggcccct tcaaaccctg tgaacacctg 1020  
 cttgatggct ggctgatcag aattggagtg tggaccatag cagtctctggc acttacttgt 1080  
 aatgcttttg tgacttcaac agttttcaga tcccctctgt acatttcccc cattaaactg 1140  
 ttaattgggg tcatcgcagc agtgaacatg ctacacggag tctccagtgc cgtgctggct 1200  
 ggtgtggatg cgttcacttt tggcagcttt gcacgacatg gtgcctgggt ggagaatggg 1260  
 gttggttgcc atgtcattgg tttttgtcc atttttgctt cagaatcatc tgttttcctg 1320  
 ctactcttg cagccctgga gcgtgggttc tctgtgaaat attctgcaaa atttgaaacg 1380  
 aaagctccat tttctagcct gaaagtaatc attttgctct gtgcctgct ggccttgacc 1440  
 atggccgcag ttccctgct ggtgggcagc aagtatggcg cctccccctc ctgectgcct 1500  
 ttgccttttg gggagcccag caccatgggc tacatggtcg ctctcatctt gctcaattcc 1560  
 ctttgcttcc tcatgatgac cattgcctac accaagctct actgcaattt ggacaaggga 1620  
 gacctggaga atatttggga ctgctctatg gtaaaacaca ttgccctggt gctcttcacc 1680

```

aactgcatcc taaactgccc tgtggtttc ttgtccttct cctctttaat aaaccttaca 1740
tttatcagtc ctgaagtaat taagtttatc cttctggtgg tagtccact tectgcatgt 1800
ctcaatcccc ttctctacat cttgttcaat cctcacttta aggaggatct ggtgagcctg 1860
agaaaacaaa cctacgtctg gacaagatca aaacacccaa gcttgatgtc aattaactct 1920
gatgatgtcg aaaaacagtc ctgtgactca actcaagcct tggtaacctt taccagctcc 1980
agcatcactt atgacctgcc tcccagttcc gtgccatcac cagcttatcc agtgactgag 2040
agctgccatc ttctctctgt ggcatttgtc ccatgtctct aa 2082

```

<210> 4  
 <211> 693  
 <212> PRT  
 <213> human

<400> 4

Leu	His	Leu	His	Asn	Asn	Arg	Ile	His	Ser	Leu	Gly	Lys	Lys	Cys	Phe
1				5					10					15	
Asp	Gly	Leu	His	Ser	Leu	Glu	Thr	Leu	Asp	Leu	Asn	Tyr	Asn	Asn	Leu
			20					25					30		
Asp	Glu	Phe	Pro	Thr	Ala	Ile	Arg	Thr	Leu	Ser	Asn	Leu	Lys	Glu	Leu
		35					40					45			
Gly	Phe	His	Ser	Asn	Asn	Ile	Arg	Ser	Ile	Pro	Glu	Lys	Ala	Phe	Val
50					55					60					
Gly	Asn	Pro	Ser	Leu	Ile	Thr	Ile	His	Phe	Tyr	Asp	Asn	Pro	Ile	Gln
65					70				75					80	
Phe	Val	Gly	Arg	Ser	Ala	Phe	Gln	His	Leu	Pro	Glu	Leu	Arg	Thr	Leu
			85					90					95		
Thr	Leu	Asn	Gly	Ala	Ser	Gln	Ile	Thr	Glu	Phe	Pro	Asp	Leu	Thr	Gly
			100					105					110		
Thr	Ala	Asn	Leu	Glu	Ser	Leu	Thr	Leu	Thr	Gly	Ala	Gln	Ile	Ser	Ser
		115						120					125		
Leu	Pro	Gln	Thr	Val	Cys	Asn	Gln	Leu	Pro	Asn	Leu	Gln	Val	Leu	Asp
		130				135					140				
Leu	Ser	Tyr	Asn	Leu	Leu	Glu	Asp	Leu	Pro	Ser	Phe	Ser	Val	Cys	Gln
145					150				155					160	
Lys	Leu	Gln	Lys	Ile	Asp	Leu	Arg	His	Asn	Glu	Ile	Tyr	Glu	Ile	Lys
			165						170					175	
Val	Asp	Thr	Phe	Gln	Gln	Leu	Leu	Ser	Leu	Arg	Ser	Leu	Asn	Leu	Ala
			180					185					190		
Trp	Asn	Lys	Ile	Ala	Ile	Ile	His	Pro	Asn	Ala	Phe	Ser	Thr	Leu	Pro
		195					200					205			
Ser	Leu	Ile	Lys	Leu	Asp	Leu	Ser	Ser	Asn	Leu	Leu	Ser	Ser	Phe	Pro
		210				215						220			
Ile	Thr	Gly	Leu	His	Gly	Leu	Thr	His	Leu	Lys	Leu	Thr	Gly	Asn	His
225					230					235				240	
Ala	Leu	Gln	Ser	Trp	Ile	Ser	Ser	Glu	Asn	Phe	Pro	Glu	Leu	Lys	Val
			245						250					255	
Ile	Glu	Met	Pro	Tyr	Ala	Tyr	Gln	Cys	Cys	Ala	Phe	Gly	Val	Cys	Glu
			260					265					270		
Asn	Ala	Tyr	Lys	Ile	Ser	Asn	Gln	Trp	Asn	Lys	Gly	Asp	Asn	Ser	Ser
		275					280					285			
Met	Asp	Asp	Leu	His	Lys	Lys	Asp	Ala	Gly	Met	Phe	Gln	Ala	Gln	Asp
		290				295					300				
Glu	Arg	Asp	Leu	Glu	Asp	Phe	Leu	Leu	Asp	Phe	Glu	Glu	Asp	Leu	Lys
305					310					315				320	
Ala	Leu	His	Ser	Val	Gln	Cys	Ser	Pro	Ser	Pro	Gly	Pro	Phe	Lys	Pro
			325						330					335	
Cys	Glu	His	Leu	Leu	Asp	Gly	Trp	Leu	Ile	Arg	Ile	Gly	Val	Trp	Thr
			340					345					350		

Ile Ala Val Leu Ala Leu Thr Cys Asn Ala Leu Val Thr Ser Thr Val  
 355 360 365  
 Phe Arg Ser Pro Leu Tyr Ile Ser Pro Ile Lys Leu Leu Ile Gly Val  
 370 375 380  
 Ile Ala Ala Val Asn Met Leu Thr Gly Val Ser Ser Ala Val Leu Ala  
 385 390 395 400  
 Gly Val Asp Ala Phe Thr Phe Gly Ser Phe Ala Arg His Gly Ala Trp  
 405 410 415  
 Trp Glu Asn Gly Val Gly Cys His Val Ile Gly Phe Leu Ser Ile Phe  
 420 425 430  
 Ala Ser Glu Ser Ser Val Phe Leu Leu Thr Leu Ala Ala Leu Glu Arg  
 435 440 445  
 Gly Phe Ser Val Lys Tyr Ser Ala Lys Phe Glu Thr Lys Ala Pro Phe  
 450 455 460  
 Ser Ser Leu Lys Val Ile Ile Leu Leu Cys Ala Leu Leu Ala Leu Thr  
 465 470 475 480  
 Met Ala Ala Val Pro Leu Leu Gly Gly Ser Lys Tyr Gly Ala Ser Pro  
 485 490 495  
 Leu Cys Leu Pro Leu Pro Phe Gly Glu Pro Ser Thr Met Gly Tyr Met  
 500 505 510  
 Val Ala Leu Ile Leu Leu Asn Ser Leu Cys Phe Leu Met Met Thr Ile  
 515 520 525  
 Ala Tyr Thr Lys Leu Tyr Cys Asn Leu Asp Lys Gly Asp Leu Glu Asn  
 530 535 540  
 Ile Trp Asp Cys Ser Met Val Lys His Ile Ala Leu Leu Phe Thr  
 545 550 555 560  
 Asn Cys Ile Leu Asn Cys Pro Val Ala Phe Leu Ser Phe Ser Ser Leu  
 565 570 575  
 Ile Asn Leu Thr Phe Ile Ser Pro Glu Val Ile Lys Phe Ile Leu Leu  
 580 585 590  
 Val Val Val Pro Leu Pro Ala Cys Leu Asn Pro Leu Leu Tyr Ile Leu  
 595 600 605  
 Phe Asn Pro His Phe Lys Glu Asp Leu Val Ser Leu Arg Lys Gln Thr  
 610 615 620  
 Tyr Val Trp Thr Arg Ser Lys His Pro Ser Leu Met Ser Ile Asn Ser  
 625 630 635 640  
 Asp Asp Val Glu Lys Gln Ser Cys Asp Ser Thr Gln Ala Leu Val Thr  
 645 650 655  
 Phe Thr Ser Ser Ser Ile Thr Tyr Asp Leu Pro Pro Ser Ser Val Pro  
 660 665 670  
 Ser Pro Ala Tyr Pro Val Thr Glu Ser Cys His Leu Ser Ser Val Ala  
 675 680 685  
 Phe Val Pro Cys Leu  
 690

<210> 5  
 <211> 2467  
 <212> DNA  
 <213> human

<400> 5  
 gaaaggagga aagaaaaaaa gaggaatgga aagagacaga gaaaggaaat gggagtggaa 60  
 ggaggaggga ctgctttgta actgctaaga ttgcagacag aaatagcaca caaccactgt 120  
 gagctgtatg cgattcagaa accaagacca aattttgctc actttcatta atcagttgct 180  
 cagatagaag gaaatgacat ctggttctgt cttcttctac atcttaattt ttggaaaata 240  
 tttttctcat ggggtgggac aggatgtcaa gtgctccctt ggctatttcc cctgtgggaa 300  
 catcacaaag tgcttgccctc agctcctgca ctgtaacggt gtggacgact gcgggaatca 360  
 ggccgatgag gacaactgtg gagacaacaa tggatggtcc atgcaatttg acaaatattt 420  
 tgccagttac tacaaaatga cttcccaata tccttttgag gcagaaacac ctgaatgttt 480

```

ggtcggttct gtgccagtgc aatgtctttg ccaaggctctg gagcttgact gtgatgaaac 540
caattttacga gctgttccat cggtttcttc aaatgtgact gcaatgtcac ttcagtgga 600
cttaataaga aagcttcctc ctgattgctt caagaattat catgatcttc agaagctgta 660
cctgcaaaac aataagatta catccatctc catctatgct ttcagaggac tgaatagcct 720
tactaaactg tatctcagtc ataacagaat aaccttcctg aagccgggtg tttttgaaga 780
tcttcacaga ctagaatggc tgataattga agataatcac ctcagtcgaa tttccccacc 840
aacattttat ggactaaatt ctcttattct cttagtcctg atgaataacg tcctcaccgg 900
tttacctgat aaacctctct gtcaacacat gccaaagacta cattggctgg accttgaagg 960
caaccatata cataatttaa gaaatttgac ttttatttcc tgcagtaatt taactgtttt 1020
agtgatgagg aaaaacaaaa ttaatcactt aaatgaaaat acttttgcac ctctccagaa 1080
actggatgaa ttggatttag gaagtaataa gattgaaaat cttccaccgc ttatattcaa 1140
ggacctgaag gagctgtcac aattgaatct ttcctataat ccaatccaga aaattcaagc 1200
aaaccaattt gattatcttg tcaaaactcaa gtctctcagc ctagaaggga ttgaaatttc 1260
aaatatccaa caaaggatgt tttagacctc tatgaatctc tctcacatat attttaagaa 1320
attccagtac tgtgggtatg caccacatgt tgcgagctgt aaaccaaaca ctgatggaat 1380
ttcatctcta gagaatctct tggcaagcat tattcagaga gtatttgtct gggttgatc 1440
tgcagttacc tgctttggaa acatttttgt catttgcatg cgaccttata tcaggtctga 1500
gaacaagctg tatgccatgt caatcatttc tctctgctgt gccgactgct taatgggaat 1560
atattttatc gtgacggag gctttgacct aaagtttcgt ggagaatata ataagcatgc 1620
gcagctgtgg atggagagta ctcatgttca gcttgtagga tctttggcca ttctgtccac 1680
agaagtatca gttttactgt taacatttct gacattggaa aaatacatct gcattgtcta 1740
tcctttttaga tgtgtgagac ctggaaaatg cagaacaatt acagttctga ttctcatttg 1800
gattactggt tttatagtgg ctttcattcc attgagcaat aaggaatttt tcaaaaacta 1860
ctatggcacc aatggagtat gcttccctct tcattcagaa gatacagaaa gtattggagc 1920
ccagatttat tcagtggaac tttttcttgg tattaatttg gccgcattta tcatcatagt 1980
tttttcctat ggaagcatgt tttatagtgt tcatacaagt gccataacag caactgaaat 2040
acggaatcaa gttaaaaaag agatgatcct tgccaaaacgt tttttcttta tagtatttac 2100
tgatgcatta tgctggatac ccatttttgt agtgaaaatt ctttcaactgc ttcaggtaga 2160
aataccaggt accataacct cttgggtagt gatttttatt ctgccatta acagtgcctt 2220
gaacccaatt ctctatactc tgaccacaag accattttaa gaaatgattc atcggttttg 2280
gtataactac agacaaaaga aatctatgga cagcaaaggc cagaaaacat atgctccatc 2340
attcatctgg gtggaaatgt ggccactgca ggagatgcca cctgagttaa tgaagccgga 2400
ccttttcaca taccctgtg aaatgtcact gatttctcaa tcaacgagac tcaattccta 2460
ttcatga

```

&lt;210&gt; 6

&lt;211&gt; 757

&lt;212&gt; PRT

&lt;213&gt; human

&lt;400&gt; 6

```

Met Thr Ser Gly Val Phe Phe Tyr Ile Leu Ile Phe Gly Lys Tyr
1      5      10      15
Phe Ser His Gly Gly Gln Asp Val Lys Cys Ser Leu Gly Tyr Phe
20     25     30
Pro Cys Gly Asn Ile Thr Lys Cys Leu Pro Gln Leu Leu His Cys Asn
35     40     45
Gly Val Asp Asp Cys Gly Asn Gln Ala Asp Glu Asp Asn Cys Gly Asp
50     55     60
Asn Asn Gly Trp Ser Met Gln Phe Asp Lys Tyr Phe Ala Ser Tyr Tyr
65     70     75     80
Lys Met Thr Ser Gln Tyr Pro Phe Glu Ala Glu Thr Pro Glu Cys Leu
85     90     95
Val Gly Ser Val Pro Val Gln Cys Leu Cys Gln Gly Leu Glu Leu Asp
100    105    110
Cys Asp Glu Thr Asn Leu Arg Ala Val Pro Ser Val Ser Ser Asn Val
115    120    125
Thr Ala Met Ser Leu Gln Trp Asn Leu Ile Arg Lys Leu Pro Pro Asp
130    135    140

```

Cys Phe Lys Asn Tyr His Asp Leu Gln Lys Leu Tyr Leu Gln Asn Asn  
 145 150 155 160  
 Lys Ile Thr Ser Ile Ser Ile Tyr Ala Phe Arg Gly Leu Asn Ser Leu  
 165 170 175  
 Thr Lys Leu Tyr Leu Ser His Asn Arg Ile Thr Phe Leu Lys Pro Gly  
 180 185 190  
 Val Phe Glu Asp Leu His Arg Leu Glu Trp Leu Ile Ile Glu Asp Asn  
 195 200 205  
 His Leu Ser Arg Ile Ser Pro Thr Phe Tyr Gly Leu Asn Ser Leu  
 210 215 220  
 Ile Leu Leu Val Leu Met Asn Asn Val Leu Thr Arg Leu Pro Asp Lys  
 225 230 235 240  
 Pro Leu Cys Gln His Met Pro Arg Leu His Trp Leu Asp Leu Glu Gly  
 245 250 255  
 Asn His Ile His Asn Leu Arg Asn Leu Thr Phe Ile Ser Cys Ser Asn  
 260 265 270  
 Leu Thr Val Leu Val Met Arg Lys Asn Lys Ile Asn His Leu Asn Glu  
 275 280 285  
 Asn Thr Phe Ala Pro Leu Gln Lys Leu Asp Glu Leu Asp Leu Gly Ser  
 290 295 300  
 Asn Lys Ile Glu Asn Leu Pro Pro Leu Ile Phe Lys Asp Leu Lys Glu  
 305 310 315 320  
 Leu Ser Gln Leu Asn Leu Ser Tyr Asn Pro Ile Gln Lys Ile Gln Ala  
 325 330 335  
 Asn Gln Phe Asp Tyr Leu Val Lys Leu Lys Ser Leu Ser Leu Glu Gly  
 340 345 350  
 Ile Glu Ile Ser Asn Ile Gln Gln Arg Met Phe Arg Pro Leu Met Asn  
 355 360 365  
 Leu Ser His Ile Tyr Phe Lys Lys Phe Gln Tyr Cys Gly Tyr Ala Pro  
 370 375 380  
 His Val Arg Ser Cys Lys Pro Asn Thr Asp Gly Ile Ser Ser Leu Glu  
 385 390 395 400  
 Asn Leu Leu Ala Ser Ile Ile Gln Arg Val Phe Val Trp Val Val Ser  
 405 410 415  
 Ala Val Thr Cys Phe Gly Asn Ile Phe Val Ile Cys Met Arg Pro Tyr  
 420 425 430  
 Ile Arg Ser Glu Asn Lys Leu Tyr Ala Met Ser Ile Ile Ser Leu Cys  
 435 440 445  
 Cys Ala Asp Cys Leu Met Gly Ile Tyr Leu Phe Val Ile Gly Gly Phe  
 450 455 460  
 Asp Leu Lys Phe Arg Gly Glu Tyr Asn Lys His Ala Gln Leu Trp Met  
 465 470 475 480  
 Glu Ser Thr His Cys Gln Leu Val Gly Ser Leu Ala Ile Leu Ser Thr  
 485 490 495  
 Glu Val Ser Val Leu Leu Leu Thr Phe Leu Thr Leu Glu Lys Tyr Ile  
 500 505 510  
 Cys Ile Val Tyr Pro Phe Arg Cys Val Arg Pro Gly Lys Cys Arg Thr  
 515 520 525  
 Ile Thr Val Leu Ile Leu Ile Trp Ile Thr Gly Phe Ile Val Ala Phe  
 530 535 540  
 Ile Pro Leu Ser Asn Lys Glu Phe Phe Lys Asn Tyr Tyr Gly Thr Asn  
 545 550 555 560  
 Gly Val Cys Phe Pro Leu His Ser Glu Asp Thr Glu Ser Ile Gly Ala  
 565 570 575  
 Gln Ile Tyr Ser Val Ala Ile Phe Leu Gly Ile Asn Leu Ala Ala Phe  
 580 585 590  
 Ile Ile Ile Val Phe Ser Tyr Gly Ser Met Phe Tyr Ser Val His Gln  
 595 600 605

Ser Ala Ile Thr Ala Thr Glu Ile Arg Asn Gln Val Lys Lys Glu Met  
 610 615 620  
 Ile Leu Ala Lys Arg Phe Phe Phe Ile Val Phe Thr Asp Ala Leu Cys  
 625 630 635 640  
 Trp Ile Pro Ile Phe Val Val Lys Phe Leu Ser Leu Leu Gln Val Glu  
 645 650 655  
 Ile Pro Gly Thr Ile Thr Ser Trp Val Val Ile Phe Ile Leu Pro Ile  
 660 665 670  
 Asn Ser Ala Leu Asn Pro Ile Leu Tyr Thr Leu Thr Thr Arg Pro Phe  
 675 680 685  
 Lys Glu Met Ile His Arg Phe Trp Tyr Asn Tyr Arg Gln Arg Lys Ser  
 690 695 700  
 Met Asp Ser Lys Gly Gln Lys Thr Tyr Ala Pro Ser Phe Ile Trp Val  
 705 710 715 720  
 Glu Met Trp Pro Leu Gln Glu Met Pro Pro Glu Leu Met Lys Pro Asp  
 725 730 735  
 Leu Phe Thr Tyr Pro Cys Glu Met Ser Leu Ile Ser Gln Ser Thr Arg  
 740 745 750  
 Leu Asn Ser Tyr Ser  
 755

&lt;210&gt; 7

&lt;211&gt; 3584

&lt;212&gt; DNA

&lt;213&gt; human

&lt;400&gt; 7

ctgctttgta	actgctaaga	ttgcagacag	aaatagcaca	caaccactgt	gagctgtatg	60
cgattcagaa	accaagacca	aattttgctc	actttcatta	atcagttgct	cagatagaag	120
gaaatgacat	ctggttctgt	cttcttctac	atcttaattt	ttggaaaata	tttttctcat	180
gggggtggac	aggatgtcaa	gtgctccctt	ggctatttcc	cctgtgggaa	catcacaaag	240
tgcttgcttc	agctcctgca	ctgtaacggt	gtggacgact	gcgggaatca	ggccgatgag	300
gacaactgtg	tggtggtttt	gtgccagtgc	atgtctttgc	caggctctga	gcttgactgg	360
atgaaaccat	ttacgagtgt	tccatcggtt	tcttcaaattg	tgactgcaat	gtcacttcag	420
tggaacttaa	taagaaaagct	tcctcctgat	tgcttcaaga	attatcatga	tcttcagaag	480
ctggacctgc	aaaacaataa	gattacatcc	atctccatct	atgcttttcag	aggactgaat	540
agccttacta	aactgtatct	cagtcataac	agaataacct	tctgaagcc	gggtgttttt	600
gaagatcttc	acagactaga	atggctgata	attgaagata	atcacctcag	tcgaatttcc	660
ccaccaacat	tttatggact	aaattctctt	attctcttag	tcctgatgaa	taacgtcttc	720
acccgtttac	ctgataaaacc	tctctgtcaa	cacatgccaa	gactacattg	gctggacctt	780
gaaggcaacc	atatccataa	tttaagaaat	ttgactttta	tttctgcag	taatttaact	840
gttttagtga	tgaggaaaaa	caaaattaat	cacttaaatg	aaaatacttt	tgacactctc	900
cagaaactgg	atgaattgga	tttaggaagt	aataagattg	aaaatcttcc	accgcttata	960
ttcaaggacc	tgaaggagct	gtcacaattg	aatctttcct	ataatccaat	ccagaaaatt	1020
caagcaaac	aatttgatta	tcttgtaaaa	ctcaagtctc	tcagcctaga	agggattgaa	1080
atttcaaata	tccaacaaag	gatgtttaga	cctcttatga	atctctctca	catatatatt	1140
aagaaattcc	agtactgtgg	gtatgcacca	catgttcgca	gctgtaaaacc	aaacactgat	1200
ggaatttcat	ctctagagaa	tctcttggca	agcattattc	agagagtatt	tgtctgggtt	1260
gtatctgcag	ttacctgctt	tggaaacatt	tttgtcattt	gcatgcgacc	ttatatcagg	1320
tctgagaaca	agctgtatgc	catgtcaatc	atttctctct	gctgtgccga	ctgcttaattg	1380
ggaatatatt	tattcgtgat	cggaggcttt	gacctaaaagt	ttcgtggaga	atacaataag	1440
catgcgcagc	tgtggatgga	gagtactcat	tgtcagcttg	taggatcttt	ggccattctg	1500
tccacagaag	tatcagtttt	actgttaaca	tttctgacat	tggaaaaata	catctgcatt	1560
gtctatcctt	ttagatgtgt	gagacctgga	aaatgcagaa	caattacagt	tctgattctc	1620
atttggatta	ctggttttat	agtggctttc	attccattga	gcaataagga	atttttcaaa	1680
aactactatg	gcaccaatgg	agtatgcttc	cctcttcatt	cagaagatac	agaaagtatt	1740
ggagcccaga	tttattcagt	ggcaattttt	cttgggtatta	atttggccgc	atttatcatc	1800
atagtttttt	cctatggaag	catgttttat	agtgttcac	aaagtgccat	aacagcaact	1860
gaaatacggg	atcaagttaa	aaaagagatg	atccttgcca	aacgtttttt	ctttatagta	1920



```

tttactgatg cattatgctg gataccatt tttgtagtga aatttctttc actgcttcag 1980
gtagaataac caggtaccat aacctcttgg gtagtgattt ttattctgcc cattaacagt 2040
gctttgaacc caattctcta tactctgacc acaagaccat ttaaagaaat gattcatcgg 2100
ttttggtata actacagaca aagaaaatct atggacagca aaggtcagaa aacatagct 2160
ccatcattca tctgggtgga aatgtggcca ctgcaggaga tgccacctga gttaatgaag 2220
ccggaccttt tcacataccc ctgtgaaatg tcaactgattt ctcaatcaac gagactcaat 2280
tcctattcat gactgactct gaaattcatt tcttcgcaga gaatactgtg ggggtgcttc 2340
atgagggatt tactgggtatg aaaatgaata ccacaaaatt aatttataat aatagctaag 2400
ataaatatatt tacaaggaca tgaggaaaaa taaaaatgac taatgctctt acaaaggga 2460
gtaattatat caataatgta tatatattag tagacatttt gcataagaaa ttaagagaaa 2520
tctacttcag taacattcat tcatTTTTtT aacatgcatt tattgagtac ccactactat 2580
gtgcatagca ttgcaatata gtcctggaag tagacagtgc agaacctttc aatctgtaga 2640
tagtgtttaa tgacaaaaga ctatacaaa gtcactgtca gttcctagtt taaagtagag 2700
ctttacctgt catgtgcatc agcaagaatc ataggcactt ttaaataaag gtttaaagtt 2760
ttggaatact cagtgtattt gcatcataga aaatgtctga ctgtttgcaa aataatattc 2820
tgTTTTaaga atccatctta cctctcttta agtttccata cacttgagag ccaacacaac 2880
atatttatta ctaaaaagat gctttgctag aaactcaaaa acagcacttc ttttggcact 2940
tcctgcccag ttttctcttt gctttaaatg aacatcatca tatggaattg gaataggaga 3000
gtatgagtac ggcagagaag tggatcagaa aaactagaat gaggataaac atttacatta 3060
gtggaaactc ctgaaataaa tccttgattt gtcagttaac tgattttcaa caaggatgcc 3120
aagacaaaaa ggcttttcaa caaaccgtgc tgttttaaga acagacctaa gtggtttaat 3180
tcacccactt tagatgggtg aatgttatgg tgtgtgaaat atctcagtaa agcagttaaa 3240
aggaaaaaga gctggaatgc actgattcag gaacttaatt tcaggaagga aaggtctgta 3300
tgtacacatt tcactttaag cagaaaatct ttcttcaaga aatgacttta ctttctcttt 3360
gcactgccag cacgtgagat actaactttt taactagtgt ttcttctcta gtctctacct 3420
tattagnatt ttttgctttc ataattgtaa acctttaagc aggagaagaa aatgTTTTca 3480
gatagtttca aatacnccaa aaatgtttgc aacacaaaaa tactggaatc naaccataat 3540
gcccttattg aatatatagt tgtatagntt tgttctgaaa accc 3584

```

<210> 8  
 <211> 722  
 <212> PRT  
 <213> human

<400> 8

```

Met Thr Ser Gly Ser Val Phe Phe Tyr Ile Leu Ile Phe Gly Lys Tyr
1      5      10      15
Phe Ser His Gly Gly Gln Asp Val Lys Cys Ser Leu Gly Tyr Phe
20     25     30
Pro Cys Gly Asn Ile Thr Lys Cys Leu Pro Gln Leu Leu His Cys Asn
35     40     45
Gly Val Asp Asp Cys Gly Asn Gln Ala Asp Glu Asp Asn Cys Val Val
50     55     60
Val Leu Cys Gln Cys Met Ser Leu Pro Gly Leu Glu Leu Asp Trp Met
65     70     75     80
Lys Pro Phe Thr Ser Val Pro Ser Val Ser Ser Asn Val Thr Ala Met
85     90     95
Ser Leu Gln Trp Asn Leu Ile Arg Lys Leu Pro Pro Asp Cys Phe Lys
100    105    110
Asn Tyr His Asp Leu Gln Lys Leu Asp Leu Gln Asn Asn Lys Ile Thr
115    120    125
Ser Ile Ser Ile Tyr Ala Phe Arg Gly Leu Asn Ser Leu Thr Lys Leu
130    135    140
Tyr Leu Ser His Asn Arg Ile Thr Phe Leu Lys Pro Gly Val Phe Glu
145    150    155    160
Asp Leu His Arg Leu Glu Trp Leu Ile Ile Glu Asp Asn His Leu Ser
165    170    175
Arg Ile Ser Pro Pro Thr Phe Tyr Gly Leu Asn Ser Leu Ile Leu Leu
180    185    190

```

Val Leu Met Asn Asn Val Leu Thr Arg Leu Pro Asp Lys Pro Leu Cys  
 195 200 205  
 Gln His Met Pro Arg Leu His Trp Leu Asp Leu Glu Gly Asn His Ile  
 210 215 220  
 His Asn Leu Arg Asn Leu Thr Phe Ile Ser Cys Ser Asn Leu Thr Val  
 225 230 235 240  
 Leu Val Met Arg Lys Asn Lys Ile Asn His Leu Asn Glu Asn Thr Phe  
 245 250 255  
 Ala Pro Leu Gln Lys Leu Asp Glu Leu Asp Leu Gly Ser Asn Lys Ile  
 260 265 270  
 Glu Asn Leu Pro Pro Leu Ile Phe Lys Asp Leu Lys Glu Leu Ser Gln  
 275 280 285  
 Leu Asn Leu Ser Tyr Asn Pro Ile Gln Lys Ile Gln Ala Asn Gln Phe  
 290 295 300  
 Asp Tyr Leu Val Lys Leu Lys Ser Leu Ser Leu Glu Gly Ile Glu Ile  
 305 310 315 320  
 Ser Asn Ile Gln Gln Arg Met Phe Arg Pro Leu Met Asn Leu Ser His  
 325 330 335  
 Ile Tyr Phe Lys Lys Phe Gln Tyr Cys Gly Tyr Ala Pro His Val Arg  
 340 345 350  
 Ser Cys Lys Pro Asn Thr Asp Gly Ile Ser Ser Leu Glu Asn Leu Leu  
 355 360 365  
 Ala Ser Ile Ile Gln Arg Val Phe Val Trp Val Val Ser Ala Val Thr  
 370 375 380  
 Cys Phe Gly Asn Ile Phe Val Ile Cys Met Arg Pro Tyr Ile Arg Ser  
 385 390 395 400  
 Glu Asn Lys Leu Tyr Ala Met Ser Ile Ile Ser Leu Cys Cys Ala Asp  
 405 410 415  
 Cys Leu Met Gly Ile Tyr Leu Phe Val Ile Gly Gly Phe Asp Leu Lys  
 420 425 430  
 Phe Arg Gly Glu Tyr Asn Lys His Ala Gln Leu Trp Met Glu Ser Thr  
 435 440 445  
 His Cys Gln Leu Val Gly Ser Leu Ala Ile Leu Ser Thr Glu Val Ser  
 450 455 460  
 Val Leu Leu Leu Thr Phe Leu Thr Leu Glu Lys Tyr Ile Cys Ile Val  
 465 470 475 480  
 Tyr Pro Phe Arg Cys Val Arg Pro Gly Lys Cys Arg Thr Ile Thr Val  
 485 490 495  
 Leu Ile Leu Ile Trp Ile Thr Gly Phe Ile Val Ala Phe Ile Pro Leu  
 500 505 510  
 Ser Asn Lys Glu Phe Phe Lys Asn Tyr Tyr Gly Thr Asn Gly Val Cys  
 515 520 525  
 Phe Pro Leu His Ser Glu Asp Thr Glu Ser Ile Gly Ala Gln Ile Tyr  
 530 535 540  
 Ser Val Ala Ile Phe Leu Gly Ile Asn Leu Ala Ala Phe Ile Ile Ile  
 545 550 555 560  
 Val Phe Ser Tyr Gly Ser Met Phe Tyr Ser Val His Gln Ser Ala Ile  
 565 570 575  
 Thr Ala Thr Glu Ile Arg Asn Gln Val Lys Lys Glu Met Ile Leu Ala  
 580 585 590  
 Lys Arg Phe Phe Phe Ile Val Phe Thr Asp Ala Leu Cys Trp Ile Pro  
 595 600 605  
 Ile Phe Val Val Lys Phe Leu Ser Leu Leu Gln Val Glu Ile Pro Gly  
 610 615 620  
 Thr Ile Thr Ser Trp Val Val Ile Phe Ile Leu Pro Ile Asn Ser Ala  
 625 630 635 640  
 Leu Asn Pro Ile Leu Tyr Thr Leu Thr Thr Arg Pro Phe Lys Glu Met  
 645 650 655

Ile	His	Arg	Phe	Trp	Tyr	Asn	Tyr	Arg	Gln	Arg	Lys	Ser	Met	Asp	Ser
			660					665					670		
Lys	Gly	Gln	Lys	Thr	Tyr	Ala	Pro	Ser	Phe	Ile	Trp	Val	Glu	Met	Trp
		675					680					685			
Pro	Leu	Gln	Glu	Met	Pro	Pro	Glu	Leu	Met	Lys	Pro	Asp	Leu	Phe	Thr
	690					695					700				
Tyr	Pro	Cys	Glu	Met	Ser	Leu	Ile	Ser	Gln	Ser	Thr	Arg	Leu	Asn	Ser
705					710					715					720
Tyr	Ser														

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/06573

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07K 14/705; C12N 15/12, 15/63, 15/70, 15/79

US CL : 530/350; 435/69.1, 252.3, 254.11, 320.1, 325

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/350; 435/69.1, 252.3, 254.11, 320.1, 325

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, Biosis, Medline, WPI

search terms: G-protein coupled receptor, Leucine rich repeats, Gonadotropin receptor, Thyrotropin receptor

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,614,363 A (CONE) 25 March 1997, entire document.	1-11
X, P — Y, P	US 5,858,716 A (ELSHOURBAGY et al.) 12 January 1999, columns 20-30, entire document.	11 — 1-10
X — Y	HWANG et al. Analysis of Expressed Sequence Tags from a Fetal Human Heart cDNA Library. Genomics. 1995, Vol. 30, pages 293-298, entire document.	5, 7, 11 — 1-4, 6, 8-10



Further documents are listed in the continuation of Box C.



See patent family annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*B\* earlier document published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\*

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\*

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\*

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*A\*

document member of the same patent family

Date of the actual completion of the international search

11 JUNE 1999

Date of mailing of the international search report

02 AUG 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Sally P. Teng

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)\*

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/06573

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-11, drawn to nucleic acids encoding LGR4, the LGR4 polypeptide, and method of using the LGR4 nucleic acid.

Group II, claims 1-11, drawn to nucleic acids encoding LGR5, the LGR5 polypeptide and method of using the LGR5 nucleic acid.

Group III, claims 1-11, drawn to nucleic acid encoding LGR6, the LGR6 polypeptide, and method of using the LGR6 nucleic acid.

Group IV, claims 12 and 13, drawn to antibody that binds to LGR4.

Group V, claims 12 and 13, drawn to antibody that binds to LGR5.

Group VI, claims 12 and 13, drawn to antibody that binds to LGR7.

Group VII, claims 14-17, drawn to transgenic animal model containing an altered LGR4 gene.

Group VIII, claims 14-17, drawn to transgenic animal model containing an altered LGR5 gene.

Group IX, claims 14-17, drawn to transgenic animal model containing an altered LGR7 gene.

Group X, claim 18, drawn to a method of screening for a ligand for LGR4.

Group XI, claim 18, drawn to a method of screening for a ligand for LGR5.

Group XII, claim 18, drawn to a method of screening for a ligand for LGR7.

Each of the claims 1-18 is in three different groups because LGR4, LGR5, and LGR7 are structurally and functionally distinct polypeptides.

The inventions listed as Groups I-XII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The special technical feature of Group I is the nucleic acid sequence encoding LGR4. The special technical feature of Group II is the nucleic acid sequence encoding LGR5. The special technical feature of Group III is the nucleic acid sequence encoding LGR7. The special technical feature of Group IV is the antibody that binds to LGR4 but does not have the amino acid sequence of LGR4. The special technical feature of Group V is the antibody that binds to LGR5 but does not have the amino acid sequence of LGR5. The special technical feature of Group VI is the antibody that binds to LGR6 but does not have the amino acid sequence of LGR6. The special technical feature of Group VII is a transgenic animal containing an altered LGR4 gene. The special technical feature of Group VIII is a transgenic animal containing an altered LGR5 gene. The special technical feature of Group IX is a transgenic animal containing an altered LGR7 gene. The special technical feature of Group X is a method of screening for a ligand that binds LGR4. The special technical feature of Group XI is a method of screening for a ligand that binds LGR5. The special technical feature of Group XII is a method of screening for a ligand that binds LGR7. The special technical feature of each group is not the same or does not correspond to the special technical feature of any other group because the products of Groups I-IX are structurally and functionally distinct and the methods of Groups I-III and X-XII are distinct methods of using different starting reagent for accomplishing different goals. The groups are not linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/06573

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	HSU et al. Characterization of Two LGR Genes Homologous to Gonadotropin and Thyrotropin Receptors with Extracellular Leucine-Rich Repeats and a G Protein-Coupled, Seven Transmembrane Region. Molecular Endocrinology. December 1998, Vol. 12, No. 12, pages 1830-1845, especially pages 1831-1837.	1-11

Form PCT/ISA/210 (continuation of second sheet)(July 1992)\*